



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

|  |                  |  |
|--|------------------|--|
| <p>(51) International Patent Classification 6 :<br/> <b>C07C 311/29, C07D 295/18, 213/56,<br/> A61K 31/535, 31/44, 31/495, 31/40, 31/18</b></p>  | <p><b>A1</b></p> | <p>(11) International Publication Number: <b>WO 96/27583</b></p> <p>(43) International Publication Date: <b>12 September 1996 (12.09.96)</b></p>   |
| <p>(21) International Application Number: <b>PCT/US96/02679</b></p> <p>(22) International Filing Date: <b>7 March 1996 (07.03.96)</b></p> <p>(30) Priority Data:<br/> <b>08/401,049                      8 March 1995 (08.03.95)                      US</b></p> <p>(60) Parent Application or Grant<br/> (63) Related by Continuation<br/> <b>US    08/401,049 (CON)</b><br/> <b>Filed on                                      8 March 1995 (08.03.95)</b></p> <p>(71) Applicant (for all designated States except US): <b>PFIZER INC.</b><br/> <b>[US/US]; 235 East 42nd Street, New York, NY 10017-5755</b><br/> <b>(US).</b></p> <p>(72) Inventors; and<br/> (75) Inventors/Applicants (for US only): <b>ROBINSON, Ralph, P.</b><br/> <b>[US/US]; 30 Friartuck Drive, Gales Ferry, CT 06335 (US).</b><br/> <b>RIZZI, James, P. [US/US]; 34 Devonshire Drive, Waterford,</b><br/> <b>CT 06335 (US).</b></p> |                  | <p>(81) Designated States: <b>AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</b></p> <p><b>Published</b><br/> <i>With international search report.</i></p> |
| <p>(54) Title: <b>ARYLSULFONYLAMINO HYDROXAMIC ACID DERIVATIVES</b></p>  |                  |  |
| <p>(57) Abstract</p> <p>A compound of formula (I), wherein n, X, R<sup>3</sup>, R<sup>4</sup> and Ar are as defined above, useful in the treatment of a condition selected from the group consisting of arthritis, cancer, tissue ulceration, restenosis, periodontal disease, epidermolysis bullosa, scleritis and other diseases characterized by matrix metalloproteinase activity, AIDS, sepsis, septic shock and other diseases involving the production of TNF.</p> <div style="text-align: center;"> <p style="text-align: right;">(I)</p> </div>   |                  |  |

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

|    |                          |    |  |    |                          |
|----|--------------------------|----|--|----|--------------------------|
| AM | Armenia                  | GB | United Kingdom                           | MW | Malawi                   |
| AT | Austria                  | GE | Georgia                                  | MX | Mexico                   |
| AU | Australia                | GN | Guinea                                   | NE | Niger                    |
| BB | Barbados                 | GR | Greece                                   | NL | Netherlands              |
| BE | Belgium                  | HU | Hungary                                  | NO | Norway                   |
| BF | Burkina Faso             | IE | Ireland                                  | NZ | New Zealand              |
| BG | Bulgaria                 | IT | Italy                                    | PL | Poland                   |
| BJ | Benin                    | JP | Japan                                    | PT | Portugal                 |
| BR | Brazil                   | KE | Kenya                                    | RO | Romania                  |
| BY | Belarus                  | KG | Kyrgyzstan                               | RU | Russian Federation       |
| CA | Canada                   | KP | Democratic People's Republic<br>of Korea | SD | Sudan                    |
| CF | Central African Republic | KR | Republic of Korea                        | SE | Sweden                   |
| CG | Congo                    | KZ | Kazakhstan                               | SG | Singapore                |
| CH | Switzerland              | LI | Liechtenstein                            | SI | Slovenia                 |
| CI | Côte d'Ivoire            | LK | Sri Lanka                                | SK | Slovakia                 |
| CM | Cameroon                 | LR | Liberia                                  | SN | Senegal                  |
| CN | China                    | LT | Lithuania                                | SZ | Swaziland                |
| CS | Czechoslovakia           | LU | Luxembourg                               | TD | Chad                     |
| CZ | Czech Republic           | LV | Latvia                                   | TG | Togo                     |
| DE | Germany                  | MC | Monaco                                   | TJ | Tajikistan               |
| DK | Denmark                  | MD | Republic of Moldova                      | TT | Trinidad and Tobago      |
| EE | Estonia                  | MG | Madagascar                               | UA | Ukraine                  |
| ES | Spain                    | ML | Mali                                     | UG | Uganda                   |
| FI | Finland                  | MN | Mongolia                                 | US | United States of America |
| FR | France                   | MR | Mauritania                               | UZ | Uzbekistan               |
| GA | Gabon                    |    |  | VN | Viet Nam                 |

- 1 -

**ARYLSULFONYLAMINO HYDROXAMIC ACID DERIVATIVES****Background of the Invention**

The present invention relates to arylsulfonfylamino hydroxamic acid derivatives which are inhibitors of matrix metalloproteinases or the production of tumor necrosis factor (hereinafter also referred to as TNF) and as such are useful in the treatment of a condition selected from the group consisting of arthritis, cancer, tissue ulceration, restenosis, periodontal disease, epidermolysis bullosa, scleritis and other diseases characterized by matrix metalloproteinase activity, AIDS, sepsis, septic shock and other diseases involving the production of TNF.

This invention also relates to a method of using such compounds in the treatment of the above diseases in mammals, especially humans, and to the pharmaceutical compositions useful therefor.

There are a number of enzymes which effect the breakdown of structural proteins and which are structurally related metalloproteases. Matrix-degrading metalloproteinases, such as gelatinase, stromelysin and collagenase, are involved in tissue matrix degradation (e.g. collagen collapse) and have been implicated in many pathological conditions involving abnormal connective tissue and basement membrane matrix metabolism, such as arthritis (e.g. osteoarthritis and rheumatoid arthritis), tissue ulceration (e.g. corneal, epidermal and gastric ulceration), abnormal wound healing, periodontal


- 2 -

disease, bone disease (e.g. Paget's disease and osteoporosis), tumor metastasis or invasion, as well as HIV-infection (J. Leuk. Biol., 52 (2):244-248, 1992).

Tumor necrosis factor is recognized to be involved in many infectious and auto-immune diseases (W. Friers, FEBS Letters, 1991, 285, 199). Furthermore, it has been shown that TNF is the prime mediator of the inflammatory response seen in sepsis and septic shock (C.E. Spooner et al., Clinical Immunology and Immunopathology, 1992, 62 S11).

#### Summary of the Invention

The present invention relates to a compound of the formula



or the pharmaceutically acceptable salts thereof, wherein


- 25                   n is 1 to 6;  
                   X is hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or NR<sup>1</sup>R<sup>2</sup> wherein R<sup>1</sup> and R<sup>2</sup> are each independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, piperidyl, (C<sub>1</sub>-C<sub>6</sub>)alkylpiperidyl, (C<sub>6</sub>-C<sub>10</sub>)arylpiperidyl, (C<sub>5</sub>-C<sub>9</sub>)heteroarylpiperidyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperidyl, (C<sub>5</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperidyl, (C<sub>1</sub>-C<sub>6</sub>)acylpiperidyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>5</sub>-C<sub>9</sub>)heteroaryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>5</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>5</sup>(C<sub>2</sub>-

- 3 -

- $C_6$ )alkyl,  $(C_1-C_5)$ alkyl( $CHR^5$ )  $(C_1-C_6)$ alkyl wherein  $R^5$  is hydroxy,  $(C_1-C_6)$ acyloxy,  $(C_1-C_6)$ alkoxy, piperazino,  $(C_1-C_6)$ alkylamino,  $(C_1-C_6)$ alkylthio,  $(C_6-C_{10})$ arylthio,  $(C_1-C_6)$ alkylsulfinyl,  $(C_6-C_{10})$ arylsulfinyl,  $(C_1-$   
 5  $C_6)$ alkylsulfoxyl,  $(C_6-C_{10})$ arylsulfoxyl, amino,  $(C_1-C_6)$ alkylamino,  $((C_1-C_6)alkyl)_2$ amino,  $(C_1-C_6)$ acylpiperazino,  $(C_1-C_6)$ alkylpiperazino,  $(C_6-C_{10})$ aryl  $(C_1-C_6)$ alkylpiperazino,  $(C_5-C_9)$ heteroaryl  $(C_1-C_6)$ alkylpiperazino, morpholino, thiomorpholino,  
 10 piperidino or pyrrolidino;  $R^6$   $(C_1-C_6)$ alkyl,  $(C_1-C_5)$ alkyl( $CHR^6$ )  $(C_1-C_6)$ alkyl wherein  $R^6$  is piperidyl,  $(C_1-C_6)$ alkylpiperidyl,  $(C_6-C_{10})$ aryl piperidyl,  $(C_6-C_{10})$ aryl  $(C_1-C_6)$ alkylpiperidyl,  $(C_5-C_9)$ heteroaryl piperidyl or  $(C_5-C_9)$ heteroaryl  $(C_1-C_6)$ alkylpiperidyl; and  $CH(R^7)COR^8$  wherein  $R^7$  is  
 15 hydrogen,  $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl  $(C_1-C_6)$ alkyl,  $(C_5-C_9)$ heteroaryl  $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylthio  $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ arylthio  $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylsulfinyl  $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ arylsulfinyl  $(C_1-C_6)$ alkyl,  
 20  $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylsulfonyl  $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ arylsulfonyl  $(C_1-C_6)$ alkyl, hydroxy  $(C_1-C_6)$ alkyl, amino  $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylamino  $(C_1-C_6)$ alkyl,  $((C_1-C_6)alkylamino)_2$   $(C_1-C_6)$ alkyl,  $R^9R^{10}NCO(C_1-C_6)alkyl$  or  $R^9OCO(C_1-C_6)alkyl$  wherein  $R^9$  and  $R^{10}$  are each  
 25 independently selected from the group consisting of hydrogen,  $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl  $(C_1-C_6)$ alkyl and  $(C_5-C_9)$ heteroaryl  $(C_1-C_6)$ alkyl; and  $R^8$  is  $R^{11}O$  or  $R^{11}R^{12}N$  wherein  $R^{11}$  and  $R^{12}$  are each independently selected from the group consisting of hydrogen,  $(C_1-$   
 30  $C_6)$ alkyl,  $(C_6-C_{10})$ aryl  $(C_1-C_6)$ alkyl and  $(C_5-C_9)$ heteroaryl  $(C_1-C_6)$ alkyl;  
 or  $R^1$  and  $R^2$ , or  $R^9$  and  $R^{10}$ , or  $R^{11}$  and  $R^{12}$  may be taken together to form an azetidiny, pyrrolidinyl, morpholinyl, thiomorpholinyl, indolinyl,  
 35 isoindolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl,  $(C_1-C_6)$ acylpiperazinyl,  $(C_1-C_6)$ alkylpiperazinyl,  $(C_6-C_{10})$ aryl piperazinyl,  $(C_5-$

- 4 -

C<sub>9</sub>)heteroarylpiperazinyl or a bridged diazabicycloalkyl ring selected from the group consisting of



wherein  $r$  is 1, 2 or 3;

$m$  is 1 or 2;

20  $p$  is 0 or 1; and

$Q$  is hydrogen, (C<sub>1</sub>-C<sub>3</sub>)alkyl or (C<sub>1</sub>-C<sub>6</sub>)acyl;

$R^3$  and  $R^4$  are each independently selected

from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl,

trifluoromethyl, trifluoromethyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-

25 C<sub>6</sub>)alkyl (difluoromethylene), (C<sub>1</sub>-

C<sub>3</sub>)alkyl(difluoromethylene)(C<sub>1</sub>-C<sub>3</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl,

(C<sub>5</sub>-C<sub>9</sub>)heteroaryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>5</sub>-

C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>6</sub>-C<sub>10</sub>)aryl,


(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-

30 C<sub>6</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl,

hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)acyloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-

C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, piperazinyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-

C<sub>6</sub>)acylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, piperidyl, (C<sub>1</sub>-C<sub>6</sub>)alkylpiperidyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>5</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylthio(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylthio(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylsulfinyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)arylsulfonyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, ((C<sub>1</sub>-C<sub>6</sub>)alkylamino)<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>13</sup>CO(C<sub>1</sub>-C<sub>6</sub>)alkyl wherein R<sup>13</sup> is R<sup>20</sup>O or R<sup>20</sup>R<sup>21</sup>N wherein R<sup>20</sup> and R<sup>21</sup> are each independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>5</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl; or R<sup>14</sup>(C<sub>1</sub>-C<sub>6</sub>)alkyl wherein R<sup>14</sup> is (C<sub>1</sub>-C<sub>6</sub>)acylpiperazino, (C<sub>6</sub>-C<sub>10</sub>)arylpiperazino, (C<sub>5</sub>-C<sub>9</sub>)heteroarylpiperazino, (C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino, (C<sub>5</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino, morpholino, thiomorpholino, piperidino, pyrrolidino, piperidyl, (C<sub>1</sub>-C<sub>6</sub>)alkylpiperidyl, (C<sub>6</sub>-C<sub>10</sub>)arylpiperidyl, (C<sub>5</sub>-C<sub>9</sub>)heteroarylpiperidyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperidyl, (C<sub>5</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperidyl or (C<sub>1</sub>-C<sub>6</sub>)acylpiperidyl; or R<sup>3</sup> and R<sup>4</sup>, or R<sup>20</sup> and R<sup>21</sup> may be taken together to form a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, oxacyclohexyl, thiocyclohexyl, indanyl or tetralinyl ring or a group of the formula



wherein R<sup>15</sup> is hydrogen, (C<sub>1</sub>-C<sub>6</sub>)acyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl,  
35 (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>5</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl  
or (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl; and  
Ar is (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>5</sub>-C<sub>9</sub>)heteroaryl, (C<sub>1</sub>-  
C<sub>6</sub>)alkyl(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>6</sub>-C<sub>10</sub>)aryl, ((C<sub>1</sub>-

- 6 -

- C<sub>6</sub>)alkoxy)<sub>2</sub>(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryloxy(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>5</sub>-C<sub>9</sub>)heteroaryloxy(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>1</sub>-C<sub>6</sub>)alkyl(C<sub>5</sub>-C<sub>9</sub>)heteroaryl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>5</sub>-C<sub>9</sub>)heteroaryl, ((C<sub>1</sub>-C<sub>6</sub>)alkoxy)<sub>2</sub>(C<sub>5</sub>-C<sub>9</sub>)heteroaryl, (C<sub>6</sub>-C<sub>10</sub>)aryloxy(C<sub>5</sub>-C<sub>9</sub>)heteroaryl, (C<sub>5</sub>-C<sub>9</sub>)heteroaryloxy(C<sub>5</sub>-C<sub>9</sub>)heteroaryl;
- 5       with the proviso that when either R<sup>1</sup> or R<sup>2</sup> is CH(R<sup>7</sup>)COR<sup>8</sup> wherein R<sup>7</sup> and R<sup>8</sup> are as defined above, the other of R<sup>1</sup> or R<sup>2</sup> is hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl or benzyl.

      The term "alkyl", as used herein, unless

10   otherwise indicated, includes saturated monovalent hydrocarbon radicals having straight, branched or cyclic moieties or combinations thereof.

      The term "alkoxy", as used herein, includes O-alkyl groups wherein "alkyl" is defined above.

- 15       The term "aryl", as used herein, unless otherwise indicated, includes an organic radical derived from an aromatic hydrocarbon by removal of one hydrogen, such as phenyl or naphthyl, optionally substituted by 1 to 3 substituents selected from the
- 20   group consisting of fluoro, chloro, trifluoromethyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>6</sub>-C<sub>10</sub>)aryloxy, trifluoromethoxy, difluoromethoxy and (C<sub>1</sub>-C<sub>6</sub>)alkyl.

- The term "heteroaryl", as used herein, unless otherwise indicated, includes an organic radical
- 25   derived from an aromatic heterocyclic compound by removal of one hydrogen, such as pyridyl, furyl, pyrolyl, thienyl, isothiazolyl, imidazolyl, benzimidazolyl, terazolyl, pyrazinyl, pyrimidyl, quinolyl, isoquinolyl, benzofuryl, isobenzofuryl,
- 30   benzothienyl, pyrazolyl, indolyl, isoindolyl, purinyl, carbazolyl, isoxazolyl, thiazolyl, oxazolyl, benzthiazolyl or benzoxazolyl, optionally substituted by 1 to 2 substituents selected from the group consisting of fluoro, chloro, trifluoromethyl, (C<sub>1</sub>-
- 35   C<sub>6</sub>)alkoxy, (C<sub>6</sub>-C<sub>10</sub>)aryloxy, trifluoromethoxy, difluoromethoxy and (C<sub>1</sub>-C<sub>6</sub>)alkyl.

      The term "acyl", as used herein, unless



- 7 -

otherwise indicated, includes a radical of the general formula RCO wherein R is alkyl, alkoxy, aryl, arylalkyl or aryalkyloxy and the terms "alkyl" or "aryl" are as defined above.

5           The term "acyloxy", as used herein, includes O-acyl groups wherein "acyl" is defined above.

          The compound of formula I may have chiral centers and therefore exist in different enantiomeric forms. This invention relates to all optical isomers  
10       and stereoisomers of the compounds of formula I and mixtures thereof.

          Preferred compounds of formula I include those wherein n is 2.

          Other preferred compounds of formula I  
15       include those wherein Ar is 4-methoxyphenyl or 4-phenoxyphenyl.

          Other preferred compounds of formula I include those wherein either R<sup>3</sup> or R<sup>4</sup> is not hydrogen.


          Other preferred compounds of formula I  
20       include those wherein n is 1 and either R<sup>1</sup> or R<sup>2</sup> is hydrogen.

          Other preferred compounds of formula I include those wherein X is hydroxy, Ar is 4-methoxyphenyl or 4-phenoxyphenyl and either R<sup>3</sup> or R<sup>4</sup> is  
25       not hydrogen.

          Other preferred compounds of formula I include those wherein X is alkoxy, Ar is 4-methoxyphenyl or 4-phenoxyphenyl and either R<sup>3</sup> or R<sup>4</sup> is not hydrogen.

30       Other preferred compounds of formula I include those wherein Ar is 4-methoxyphenyl or 4-phenoxyphenyl and R<sup>3</sup> and R<sup>4</sup> are taken together to form (C<sub>3</sub>-C<sub>6</sub>)cycloalkanyl, oxacyclohexanyl, thiocyclohexanyl, indanyl or a group of the formula

- 8 -



5

wherein R<sup>15</sup> is (C<sub>1</sub>-C<sub>6</sub>)acyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>5</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl.

10

More preferred compounds of formula I are those wherein n is 2, Ar is 4-methoxyphenyl or 4-phenoxyphenyl, R<sup>1</sup> and R<sup>2</sup> are taken together to form piperazinyl, (C<sub>1</sub>-C<sub>6</sub>)alkylpiperazinyl, (C<sub>6</sub>-C<sub>10</sub>)aryl piperazinyl or (C<sub>5</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-

15

C<sub>6</sub>)alkylpiperazinyl, and either R<sup>3</sup> or R<sup>4</sup> is not hydrogen or both R<sup>3</sup> and R<sup>4</sup> are not hydrogen.

More preferred compounds of formula I are those wherein n is 2, Ar is 4-methoxyphenyl or 4-phenoxyphenyl, R<sup>1</sup> is hydrogen or (C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>2</sup> is 2-pyridylmethyl, 3-pyridylmethyl or 4-pyridylmethyl, and either R<sup>3</sup> or R<sup>4</sup> is not hydrogen or both R<sup>3</sup> and R<sup>4</sup> are not hydrogen.

20

More preferred compounds of formula I are those wherein n is 1, Ar is 4-methoxyphenyl or 4-phenoxyphenyl, R<sup>1</sup> is hydrogen, R<sup>2</sup> is 2-pyridylmethyl, 3-pyridylmethyl or 4-pyridylmethyl, and either R<sup>3</sup> or R<sup>4</sup> is not hydrogen or both R<sup>3</sup> and R<sup>4</sup> are not hydrogen.

25

More preferred compounds of formula I are those wherein n is 2, Ar is 4-methoxyphenyl, R<sup>1</sup> is hydrogen or (C<sub>1</sub>-C<sub>6</sub>)alkyl and R<sup>2</sup> is R<sup>5</sup> (C<sub>2</sub>-C<sub>6</sub>)alkyl wherein R<sup>5</sup> is morpholino, thiomorpholino, piperidino, pyrrolidino, (C<sub>1</sub>-C<sub>6</sub>)acylpiperazino, (C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino, (C<sub>6</sub>-C<sub>10</sub>)arylpiperazino, (C<sub>5</sub>-C<sub>9</sub>)heteroarylpiperazino, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino or (C<sub>5</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino and either R<sup>3</sup> or R<sup>4</sup> is not hydrogen or both R<sup>3</sup> and R<sup>4</sup> are not hydrogen.

30

35

More preferred compounds of formula I are

- 9 -

those wherein n is 1, Ar is 4-methoxyphenyl or 4-phenoxyphenyl, R<sup>1</sup> is hydrogen, R<sup>2</sup> is R<sup>5</sup> (C<sub>2</sub>-C<sub>6</sub>)alkyl wherein R<sup>5</sup> is morpholino, thiomorpholino, piperidino, pyrrolidino (C<sub>1</sub>-C<sub>6</sub>)acylpiperazino, (C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino, (C<sub>6</sub>-C<sub>10</sub>)aryl piperazino, (C<sub>5</sub>-C<sub>9</sub>)heteroaryl piperazino, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino or (C<sub>5</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino and either R<sup>3</sup> or R<sup>4</sup> is not hydrogen or both R<sup>3</sup> and R<sup>4</sup> are not hydrogen.

Specific preferred compounds of formula I include the following:

2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)(3-morpholin-4-yl-3-oxopropyl)amino]-3-methylbutyramide;

2-(R)-2-[(2-Benzylcarbamoyl ethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide;

2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)(2-[(pyridin-3-ylmethyl)-carbamoyl]ethyl)amino]-3-methylbutyramide;

2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)[2-(methylpyridin-3-ylmethylcarbamoyl)ethyl]amino]-3-methylbutyramide;

4-(3-[1-(R)-1-Hydroxycarbamoyl-2-methylpropyl)(4-methoxybenzenesulfonyl)amino]propionyl)piperazine-1-carboxylic acid, tert-butyl ester;

2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)(3-oxo-3-piperazin-1-ylpropyl)amino]-3-methylbutyramide hydrochloride;

2-(R)-2-[(Benzylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]N-hydroxy-3-methylbutyramide;

2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)-[(2-morpholin-4-ylethylcarbamoyl)methyl]amino]-3-methylbutyramide; and

2-(R)-N-Hydroxy-2-[(4-

- 10 -

methoxybenzenesulfonyl) [(pyridin-3-ylmethyl) carbamoyl] methyl] amino) -3-methylbutyramide.

Other specific compounds of formula I include the following:

- 5                   2 - (R) - 3,3,3-Trifluoro-N-hydroxy-2-  
[(methoxybenzenesulfonyl) (3-morpholin-4-yl-3-oxopropyl) amino] propionamide;
- 2 - (R) - N-Hydroxy-2 - ((4-  
phenoxybenzenesulfonyl) [2 - (methylpyridin-4-ylmethylcarbamoyl) ether] amino) -3-methylbutyramide;
- 10                   4 - [4-Methoxybenzenesulfonyl) (3-morpholin-4-yl-3-oxopropyl) amino] -1-methylpiperidene-4-carboxylic acid hydroxyamide;
- 2 - (R) - N-Hydroxy-2 - ((4-  
15 methoxybenzenesulfonyl) - [3 - (4-methylpiperazin-1-yl) -3-oxopropyl] amino) -3-methylbutyramide;
- 2 - (R) - 2 - [(2-Carboxyethyl) (4-methoxybenzenesulfonyl) amino] -N-hydroxy-3-methylbutyramide;
- 20                   [(2-Carboxyethyl) (3,4-dimethoxybenzenesulfonyl) amino] -N-hydroxy-acetamide;
- 2 - (R) - 2 - [(2-Carbamoylethyl) (4-methoxybenzenesulfonyl) amino] -N-hydroxy-3-methylbutyramide;
- 25                   2 - (R) , 3 - (R) - 3,N-Dihydroxy-2 - [(4-methoxybenzenesulfonyl) (3-oxo-3-piperidin-1-ylpropyl) amino] -butyramide;
- 2 - (R) - N-Hydroxy-2 - ((4-methoxybenzenesulfonyl) [3 - (methylpyridin-3-ylmethylcarbamoyl) propyl] amino) -3-methylbutyramide;
- 30                   2 - (R) - N-Hydroxy-2 - ((4-methoxybenzenesulfonyl) [2 - (methylcarboxymethylcarbamoyl) ethyl] amino) -3-methylbutyramide;
- 35                   2 - (R) - N-Hydroxy-2 - ((4-methoxybenzenesulfonyl) - [(1-methylpiperidin-4-ylcarbamoyl) methyl] amino) -3-methylbutyramide;

- 11 -

2 - (R) - 2 - Cyclohexyl - N - hydroxy - 2 - ((4 - methoxybenzenesulfonyl) - [3 - (4 - methylpiperazin - 1 - yl) - 3 - oxopropyl] amino) - acetamide;

2 - (R) - N - Hydroxy - 2 -  
5 [(methoxybenzenesulfonyl) (3 - morpholin - 4 - yl - 3 - oxopropyl) amino] - 4 - (morpholin - 4 - yl) butyramide.

The present invention also relates to a pharmaceutical composition for (a) the treatment of a condition selected from the group consisting of  
10 arthritis, cancer, tissue ulceration, restenosis, periodontal disease, epidermolysis bullosa, scleritis and other diseases characterized by matrix metalloproteinase activity, AIDS, sepsis, septic shock and other diseases involving the production of tumor  
15 necrosis factor (TNF) or (b) the inhibition of matrix metalloproteinases or the production of tumor necrosis factor (TNF) in a mammal, including a human, comprising an amount of a compound of claim 1 or a  
pharmaceutically acceptable salt thereof, effective in  
20 such treatments and a pharmaceutically acceptable carrier.

The present invention also relates to a method for the inhibition of (a) matrix metalloproteinases or (b) the production of tumor  
25 necrosis factor (TNF) in a mammal, including a human, comprising administering to said mammal an effective amount of a compound of claim 1 or a pharmaceutically acceptable salt thereof.


The present invention also relates to a  
30 method for treating a condition selected from the group consisting of arthritis, cancer, tissue ulceration, restenosis, periodontal disease, epidermolysis bullosa, scleritis and other diseases characterized by matrix metalloproteinase activity, AIDS, sepsis, septic shock  
35 and other diseases involving the production of tumor necrosis factor (TNF) in a mammal, including a human, comprising administering to said mammal an amount of a


- 12 -

compound of claim 1 or a pharmaceutically acceptable salt thereof, effective in treating such a condition.


Detailed Description of the Invention

5     The following reaction Schemes illustrate the preparation of the compounds of the present invention. Unless otherwise indicated  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $n$  and  $Ar$  in the reaction Schemes and the discussion that follow are defined as above.

Scheme 1


Scheme 1 cont'd



Scheme 2


VI

1



VIII

2





IV

I


35

- 16 -

Scheme 3


Scheme 4

- 18 -

Scheme 4 cont'd


XIV

15



XV

30




XVI

- 19 -

In reaction 1 of Scheme 1, the amino acid compound of formula VII, wherein  $R^{16}$  is  $(C_1-C_6)$ alkyl, benzyl, allyl or tert-butyl, is converted to the corresponding compound of formula VI by reacting VII with a reactive functional derivative of an arylsulfonic acid compound, such as an arylsulfonyl chloride, in the presence of a base, such as triethylamine, and a polar solvent, such as tetrahydrofuran, dioxane, water or acetonitrile, preferably a mixture of dioxane and water. The reaction mixture is stirred, at room temperature, for time period between about 10 minutes to about 24 hours, preferably about 60 minutes.

In reaction 2 of Scheme 1, the arylsulfonyl amino compound of formula VI, wherein  $R^{16}$  is  $(C_1-C_6)$ alkyl, benzyl, allyl or tert-butyl, is converted to the corresponding compound of formula V, wherein n is 1, 3, 4, 5 or 6, by reacting VI with a reactive derivative of an alcohol of the formula



such as the chloride, bromide or iodide derivative, preferably the bromide derivative, wherein the  $R^{17}$  protecting group is  $(C_1-C_6)$ alkyl, benzyl, allyl or tert-butyl, in the presence of a base such as potassium carbonate or sodium hydride, preferably sodium hydride, and a polar solvent, such as dimethylformamide. The reaction mixture is stirred, at room temperature, for a time period between about 60 minutes to about 48 hours, preferably about 18 hours. The  $R^{17}$  protecting group is chosen such that it may be selectively removed in the presence of and without loss of the  $R^{16}$  protecting group, therefore,  $R^{17}$  cannot be the same as  $R^{16}$ . Removal of the  $R^{17}$  protecting group from the compound of formula V to give the corresponding carboxylic acid of formula IV, in reaction 3 of Scheme 1, is carried out under conditions appropriate for that particular

- 20 -

$R^{17}$  protecting group in use which will not affect the  $R^{16}$  protecting group. Such conditions include; (a) saponification where  $R^{17}$  is  $(C_1-C_6)$ alkyl and  $R^{16}$  is tert-butyl, (b) hydrogenolysis where  $R^{17}$  is benzyl and  $R^{16}$  is tert-butyl or  $(C_1-C_6)$ alkyl, (c) treatment with a strong acid such as trifluoroacetic acid or hydrochloric acid where  $R^{17}$  is tert-butyl and  $R^{16}$  is  $(C_1-C_6)$ alkyl, benzyl or allyl, or (d) treatment with tributyltinhydride and acetic acid in the presence of catalytic bis(triphenylphosphine) palladium (II) chloride where  $R^{17}$  is allyl and  $R^{16}$  is  $(C_1-C_6)$ alkyl, benzyl or tert-butyl.

In reaction 4 of Scheme 1, the carboxylic acid of formula IV is condensed with an amine,  $R^1R^2NH$ , or the salt thereof, to give the corresponding amide compound of formula III. The formation of amides from primary or secondary amines or ammonia and carboxylic acids is achieved by conversion of the carboxylic acid to an activated functional derivative which subsequently undergoes reaction with a primary or secondary amine or ammonia to form the amide. The activated functional derivative may be isolated prior to reaction with the primary or secondary amine or ammonia. Alternatively, the carboxylic acid may be treated with oxalyl chloride or thionyl chloride, neat or in an inert solvent, such as chloroform, at a temperature between about 25°C. to about 80°C., preferably about 50°C., to give the corresponding acid chloride functional derivative. The inert solvent and any remaining oxalyl chloride or thionyl chloride is then removed by evaporation under vacuum. The remaining acid chloride functional derivative is then reacted with the primary or secondary amine or ammonia in an inert solvent, such as methylene chloride, to form the amide. The preferred method for the condensation of the carboxylic acid of formula IV with an amine to provide the corresponding amide compound of

- 21 -

formula III is the treatment of IV with (benzotriazol-1-yloxy)tris(dimethylamino) phosphonium

hexafluorophosphate in the presence of a base, such as triethylamine, to provide the benzotriazol-1-oxy ester  
5 in situ which, in turn, reacts with the amine,  $R^1R^2N$ , in an inert solvent, such as methylene chloride, at room temperature to give the amide compound of formula III.

Removal of the  $R^{16}$  protecting group from the  
10 compound of formula III to give the corresponding carboxylic acid of formula II, in reaction 5 of Scheme 1, is carried out under conditions appropriate for the particular  $R^{16}$  protecting group in use. Such conditions include; (a) saponification where  $R^{16}$  is  
15 lower alkyl, (b) hydrogenolysis where  $R^{16}$  is benzyl, (c) treatment with a strong acid, such as trifluoroacetic acid or hydrochloric acid, where  $R^{16}$  is tert-butyl, or (d) treatment with tributyltinhydride and acetic acid in the presence of catalytic  
20 bis(triphenylphosphine) palladium (II) chloride where  $R^{16}$  is allyl.

In reaction 6 of Scheme 1, the carboxylic acid compound of formula II is converted to the hydroxamic acid compound of formula I by treating II  
25 with 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide and 1-hydroxybenztriazole in a polar solvent, such as dimethylformamide, followed by the addition of hydroxylamine to the reaction mixture after a time period between about 15 minutes to about 1 hour,  
30 preferably about 30 minutes. The hydroxylamine is preferably generated in situ from a salt form, such as hydroxylamine hydrochloride, in the presence of a base, such as N-methylmorpholine. Alternatively, a protected derivative of hydroxylamine or its salt form, where the  
35 hydroxyl group is protected as a tert-butyl, benzyl or allyl ether, may be used in the presence of (benzotriazol-1-yloxy)tris(dimethylamino) phosphonium

- 22 -

hexafluorophosphate and a base, such as N-methylmorpholine. Removal of the hydroxylamine protecting group is carried out by hydrogenolysis for a benzyl protecting group or treatment with a strong  
5 acid, such as trifluoroacetic acid, for a tert-butyl protecting group. The allyl protecting group may be removed by treatment with tributyltinhydride and acetic acid in the presence of catalytic  
bis(triphenylphosphine) palladium (II) chloride. N,O-  
10 bis(4-methoxybenzyl)hydroxylamine may also be used as the protected hydroxylamine derivative where deprotection is achieved using a mixture of methanesulfonic acid and trifluoroacetic acid.

In reaction 1 of Scheme 2, the  
15 arylsulfonylamino compound of formula VI, wherein  $R^{16}$  is  $(C_1-C_6)$ alkyl, benzyl or tert-butyl, is converted to the corresponding compound of formula VIII, wherein  $R^{18}$  is 2-propenyl or 3-butenyl, by reacting IX with a reactive functional derivative, such as the halide,  
20 preferably the iodide derivative, of 2-propen-1-ol when  $R^{18}$  is 2-propenyl or 3-buten-1-ol when  $R^{18}$  is 3-butenyl, in the presence of a base, such as potassium carbonate, cesium carbonate or sodium hydride, preferably sodium hydride when  $R^{18}$  is 2-propenyl or  
25 cesium carbonate when  $R^{18}$  is 3-butenyl. The reaction is stirred in a polar solvent, such as dimethylformamide, at room temperature, for a time period between about 2 hours to about 48 hours, preferably about 18 hours.

30 In reaction 2 of Scheme 2, the compound of formula VIII is converted to the carboxylic acid compound of formula IV, wherein n is 2. The compound of formula VIII, wherein  $R^{18}$  is 2-propenyl, is converted to the compound of formula IV, wherein n is  
35 2, by reacting VIII with borane-dimethylsulfide complex, followed by immediate oxidation using chromium trioxide in aqueous acetic acid. The oxidative



- 23 -

cleavage of terminal olefins to carboxylic acids can be achieved by several methods known in the art. The preferred method for the oxidative cleavage of the compound of formula VIII, wherein  $R^{18}$  is 3-butenyl, to obtain the carboxylic acid compound of formula IV is to react VIII with sodium periodate in the presence of a catalytic amount of ruthenium (III) chloride in a mixture of carbon tetrachloride, acetonitrile and water.

The compound of formula IV, wherein  $n$  is 2, is further reacted to provide the hydroxamic acid compound of formula I, wherein  $n$  is 2, according to the procedure described above in reactions 4, 5 and 6 of Scheme 1.

An alternative method for the synthesis of the hydroxamic acid compound of formula I, wherein  $n$  is 1 and  $R^3$  and  $R^4$  are both hydrogen, is shown in reaction 1 of Scheme 3, beginning with reacting iminoacetic acid or a metal or ammonium salt of iminoacetic acid of formula X with a functional derivative of an arylsulfonic acid compound, such as an arylsulfonyl chloride, at room temperature, in the presence of a suitable base, such as triethylamine, and a polar solvent such as tetrahydrofuran, dioxane, water or acetonitrile, preferably a mixture of dioxane and water, to give the corresponding dicarboxylic acid compound of formula XI.

In reaction 2 of Scheme 3, the dicarboxylic acid compound of formula XI is dehydrated to give a cyclic anhydride compound of formula XII. The formation of cyclic anhydrides by dehydration of dicarboxylic acids may be achieved by a variety of means. The preferred method for the dehydration of the dicarboxylic acid compound of formula XI to give a cyclic anhydride compound of formula XII is to treat XI with an excess of acetic anhydride at a temperature between about 25°C. to about 80°C., preferably about

60°C. Excess acetic anhydride and acetic acid, a by-product of the reaction, are removed by evaporation under reduced pressure leaving the cyclic anhydride compound of formula XII.


5 In reaction 3 of Scheme 3, the cyclic  
anhydride compound of formula XII is reacted, at room  
temperature, with an amine,  $\text{NR}^1\text{R}^2$ , or a salt of the  
amine, such as the hydrochloride, in the presence of a  
base, such as triethylamine, to give the carboxylic  
10 acid of formula II, wherein n is 1 and  $\text{R}^3$  and  $\text{R}^4$  are  
both hydrogen. Suitable solvents for the reaction are  
those that will not react with the starting materials,  
which include chloroform, methylene chloride and  
dimethylformamide, preferably methylene chloride.

15                   The compound of formula II is further reacted to give the hydroxamic acid compound of formula I, wherein n is 1 and R<sup>3</sup> and R<sup>4</sup> are both hydrogen, according to the procedure described above in reaction 6 of Scheme 1.

20 In reaction 1 of Scheme 4, the carboxylic acid compound of formula IV, wherein n is 2, is converted to the corresponding compound of formula V, wherein R<sup>19</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl or tert-butyl, by reacting IV with a compound of the formula

25  $(R^{19}O)_2CHN(CH_3)_2$   
wherein  $R^{19}$  is  $(C_1-C_6)$ alkyl or tert-butyl, in an inert solvent, such as toluene, at a temperature between about 60°C. to about 100°C., preferably about 100°C., for a time period between about 1 hour to about 3  
30 hours, preferably 2 hours. In reaction 2 of Scheme 4, the arylsulfonyl amino compound of formula VI wherein n is 1, 3, 4, 5 or 6 and  $R^{16}$  is  $(C_1-C_6)$ alkyl, benzyl, allyl or tert-butyl, is converted to the corresponding compound of formula XIII, wherein  $R^{19}$  is  $(C_1-C_6)$ alkyl  
35 or tert-butyl, by reacting VI with a reactive derivative of an alcohol of the formula

- 25 -



such as the chloride, bromide or iodide derivative, preferably the bromide derivative, wherein  $R^{19}$  is ( $C_1$ - $C_6$ )alkyl or tert-butyl, in the presence of a base such as potassium carbonate or sodium hydride, preferably sodium hydride, and a polar solvent, such as dimethylformamide. The reaction is stirred, at room temperature, for a time period between about 60 minutes to about 48 hours, preferably about 18 hours. The  $R^{16}$  protecting group, of the compounds of formulas IV and VI, is chosen such that it may be selectively removed in the presence of and without loss of the  $R^{19}$  ( $C_1$ - $C_6$ )alkyl or tert-butyl group, therefore,  $R^{16}$  cannot be the same as  $R^{19}$ . Removal of the  $R^{16}$  protecting group from the compound of formula XIII to give the corresponding carboxylic acid of formula XIV, wherein n is 1 to 6, in reaction 3 of Scheme 4, is carried out under conditions appropriate for that particular  $R^{16}$  protecting group in use which will not affect the  $R^{19}$  ( $C_1$ - $C_6$ )alkyl or tert-butyl group. Such conditions include; (a) saponification where  $R^{16}$  is ( $C_1$ - $C_6$ )alkyl and  $R^{19}$  is tert-butyl, (b) hydrogenolysis where  $R^{16}$  is benzyl and  $R^{19}$  is tert-butyl or ( $C_1$ - $C_6$ )alkyl, (c) treatment with a strong acid such as trifluoroacetic acid or hydrochloric acid where  $R^{16}$  is tert-butyl and  $R^{19}$  is ( $C_1$ - $C_6$ )alkyl, or (d) treatment with tributyltinhydride and acetic acid in the presence of catalytic bis(triphenylphosphine) palladium (II) chloride where  $R^{16}$  is allyl and  $R^{19}$  is ( $C_1$ - $C_6$ )alkyl or tert-butyl.

In reaction 4 of Scheme 4, the carboxylic acid of formula XIV is converted to the hydroxamic acid compound of formula XV, wherein n is 1 to 6, by treating XIV with 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide and 1-hydroxybenztriazole in a polar solvent, such as dimethylformamide, followed by the

- 26 -

addition of hydroxylamine to the reaction mixture after a time period between about 15 minutes to about 1 hour, preferably about 30 minutes. The hydroxylamine is preferably generated in situ from a salt form, such as hydroxylamine hydrochloride, in the presence of a base, such as N-methylmorpholine. Alternatively, a protected derivative of hydroxylamine or its salt form, where the hydroxyl group is protected as a tert-butyl, benzyl or allyl ether, may be used in the presence of (benzotriazol-1-yloxy)tris(dimethylamino) phosphonium hexafluorophosphate and a base, such as N-methylmorpholine. Removal of the hydroxylamine protecting groups is carried out by hydrogenolysis for a benzyl protecting group or treatment with a strong acid, such as trifluoroacetic acid, for a tert-butyl protecting group. The allyl protecting group may be removed by treatment with tributyltinhydride and acetic acid in the presence of catalytic bis(triphenylphosphine) palladium (II) chloride. N,O-bis(4-methoxybenzyl)hydroxylamine may also be used, when R<sup>19</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, as the protected hydroxylamine derivative where deprotection is achieved using a mixture of methanesulfonic acid and trifluoroacetic acid.

In reaction 5 of Scheme 4, the amide formula of formula XV is, if desired, converted to the corresponding carboxylic acid compound of formula XVI by (a) saponification where R<sup>19</sup> is lower alkyl or (b) treatment with a strong acid, such as trifluoroacetic acid or hydrochloric acid, where R<sup>19</sup> is tert-butyl.

Pharmaceutically acceptable salts of the acidic compounds of the invention are salts formed with bases, namely cationic salts such as alkali and alkaline earth metal salts, such as sodium, lithium, potassium, calcium, magnesium, as well as ammonium salts, such as ammonium, trimethyl-ammonium, diethylammonium, and tris-(hydroxymethyl)-

- 27 -

methyllummonium salts.

Similarly acid addition salts, such as of mineral acids, organic carboxylic and organic sulfonic acids e.g. hydrochloric acid, methanesulfonic acid, maleic acid, are also possible provided a basic group, such as pyridyl, constitutes part of the structure.

The ability of the compounds of formula I or their pharmaceutically acceptable salts (hereinafter also referred to as the compounds of the present invention) to inhibit matrix metalloproteinases or the production of tumor necrosis factor (TNF) and, consequently, demonstrate their effectiveness for treating diseases characterized by matrix metalloproteinase or the production of tumor necrosis factor is shown by the following in vitro assay tests.

#### Biological Assay

##### Inhibition of Human Collagenase (MMP-1)

Human recombinant collagenase is activated with trypsin using the following ratio: 10  $\mu$ g trypsin per 100  $\mu$ g of collagenase. The trypsin and collagenase are incubated at room temperature for 10 minutes then a five fold excess (50  $\mu$ g/10  $\mu$ g trypsin) of soybean trypsin inhibitor is added.

10 mM stock solutions of inhibitors are made up in dimethyl sulfoxide and then diluted using the following Scheme:

10 mM ---> 120  $\mu$ M ---> 12  $\mu$ M ---> 1.2  $\mu$ M ---> 0.12  $\mu$ M

Twenty-five microliters of each concentration is then added in triplicate to appropriate wells of a 96 well microfluor plate. The final concentration of inhibitor will be a 1:4 dilution after addition of enzyme and substrate. Positive controls (enzyme, no inhibitor) are set up in wells D1-D6 and blanks (no enzyme, no inhibitors) are set in wells D7-D12.

Collagenase is diluted to 400 ng/ml and 25  $\mu$ l is then added to appropriate wells of the microfluor plate. Final concentration of collagenase in the assay

- 28 -

is 100 ng/ml.

Substrate (DNP-Pro-Cha-Gly-Cys(Me)-His-Ala-Lys(NMA)-NH<sub>2</sub>) is made as a 5 mM stock in dimethyl sulfoxide and then diluted to 20 μM in assay buffer.

5 The assay is initiated by the addition of 50 μl substrate per well of the microfluor plate to give a final concentration of 10 μM.

Fluorescence readings (360 nm excitation, 460 nm emission) were taken at time 0 and then at 20 minute  
10 intervals. The assay is conducted at room temperature with a typical assay time of 3 hours.

Fluorescence vs time is then plotted for both the blank and collagenase containing samples (data from triplicate determinations is averaged). A time point  
15 that provides a good signal (the blank) and that is on a linear part of the curve (usually around 120 minutes) is chosen to determine IC<sub>50</sub> values. The zero time is used as a blank for each compound at each concentration and these values are subtracted from the 120 minute  
20 data. Data is plotted as inhibitor concentration vs % control (inhibitor fluorescence divided by fluorescence of collagenase alone x 100). IC<sub>50</sub>'s are determined from the concentration of inhibitor that gives a signal that is 50% of the control.

25 If IC<sub>50</sub>'s are reported to be <0.03 μM then the inhibitors are assayed at concentrations of 0.3 μM, 0.03 μM, 0.03 μM and 0.003 μM.

#### Inhibition of Gelatinase (MMP-2)

Inhibition of gelatinase activity is assayed  
30 using the Dnp-Pro-Cha-Gly-Cys(Me)-His-Ala-Lys(NMA)-NH<sub>2</sub> substrate (10 μM) under the same conditions as inhibition of human collagenase (MMP-1).

72kD gelatinase is activated with 1 mM APMA (p-aminophenyl mercuric acetate) for 15 hours at 4°C.  
35 and is diluted to give a final concentration in the assay of 100 mg/ml. Inhibitors are diluted as for inhibition of human collagenase (MMP-1) to give final

- 29 -

concentrations in the assay of 30  $\mu\text{M}$ , 3  $\mu\text{M}$ , 0.3  $\mu\text{M}$  and 0.03  $\mu\text{M}$ . Each concentration is done in triplicate.

Fluorescence readings (360 nm excitation, 460 emission) are taken at time zero and then at 20 minutes intervals for 4 hours.

IC<sub>50</sub>'s are determined as per inhibition of human collagenase (MMP-1). If IC<sub>50</sub>'s are reported to be less than 0.03  $\mu\text{M}$ , then the inhibitors are assayed at final concentrations of 0.3  $\mu\text{M}$ , 0.03  $\mu\text{M}$ , 0.003  $\mu\text{M}$  and 0.003  $\mu\text{M}$ .

#### Inhibition of Stromelysin Activity (MMP-3)

Inhibition of stromelysin activity is based on a modified spectrophotometric assay described by Weingarten and Feder (Weingarten, H. and Feder, J., Spectrophotometric Assay for Vertebrate Collagenase, Anal. Biochem. 147, 437-440 (1985)). Hydrolysis of the thio peptolide substrate [Ac-Pro-Leu-Gly-SCH(CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>)CO-Leu-Gly-OC<sub>2</sub>H<sub>5</sub>] yields a mercaptan fragment that can be monitored in the presence of Ellman's reagent.

Human recombinant prostromelysin is activated with trypsin using a ratio of 1  $\mu\text{l}$  of a 10 mg/ml trypsin stock per 26  $\mu\text{g}$  of stromelysin. The trypsin and stromelysin are incubated at 37°C. for 15 minutes followed by 10  $\mu\text{l}$  of 10 mg/ml soybean trypsin inhibitor for 10 minutes at 37°C. to quench trypsin activity.

Assays are conducted in a total volume of 250  $\mu\text{l}$  of assay buffer (200 mM sodium chloride, 50 mM MES, and 10 mM calcium chloride, pH 6.0) in 96-well microliter plates. Activated stromelysin is diluted in assay buffer to 25  $\mu\text{g/ml}$ . Ellman's reagent (3-Carboxy-4-nitrophenyl disulfide) is made as a 1M stock in dimethyl formamide and diluted to 5 mM in assay buffer with 50  $\mu\text{l}$  per well yielding at 1 mM final concentration.

10 mM stock solutions of inhibitors are made in dimethyl sulfoxide and diluted serially in assay

- 30 -

buffer such that addition of 50  $\mu$ L to the appropriate wells yields final concentrations of 3  $\mu$ M, 0.3  $\mu$ M, 0.003  $\mu$ M, and 0.0003  $\mu$ M. All conditions are completed in triplicate.

5           A 300 mM dimethyl sulfoxide stock solution of the peptide substrate is diluted to 15 mM in assay buffer and the assay is initiated by addition of 50  $\mu$ l to each well to give a final concentration of 3 mM substrate. Blanks consist of the peptide substrate and  
10   Ellman's reagent without the enzyme. Product formation was monitored at 405 nm with a Molecular Devices UVmax plate reader.

IC<sub>50</sub> values were determined in the same manner as for collagenase.

15                           Inhibition of MMP-13

Human recombinant MMP-13 is activated with 2mM APMA (p-aminophenyl mercuric acetate) for 1.5 hours, at 37°C. and is diluted to 400 mg/ml in assay buffer (50 mM Tris, pH 7.5, 200 mM sodium chloride, 5mM  
20   calcium chloride, 20 $\mu$ M zinc chloride, 0.02% brij). Twenty-five microliters of diluted enzyme is added per well of a 96 well microfluor plate. The enzyme is then diluted in a 1:4 ratio in the assay by the addition of inhibitor and substrate to give a final concentration  
25   in the assay of 100 mg/ml.

10 mM stock solutions of inhibitors are made up in dimethyl sulfoxide and then diluted in assay buffer as per the inhibitor dilution scheme for inhibition of human collagenase (MMP-1): Twenty-five  
30   microliters of each concentration is added in triplicate to the microfluor plate. The final concentrations in the assay are 30  $\mu$ M, 3  $\mu$ M, 0.3  $\mu$ M and 0.03  $\mu$ M.

Substrate (Dnp-Pro-Cha-Gly-Cys(Me)-His-Ala-  
35   Lys(NMA)-NH<sub>2</sub>) is prepared as for inhibition of human collagenase (MMP-1) and 50  $\mu$ l is added to each well to give a final assay concentration of 10  $\mu$ M.



- 31 -

Fluorescence readings (360 nM excitation; 450 emission) are taken at time 0 and every 5 minutes for 1 hour.

Positive controls consist of enzyme and substrate with no inhibitor and blanks consist of substrate only.

IC<sub>50</sub>'s are determined as per inhibition of human collagenase (MMP-1). If IC<sub>50</sub>'s are reported to be less than 0.03  $\mu$ M, inhibitors are then assayed at final concentrations of 0.3  $\mu$ M, 0.03  $\mu$ M, 0.003  $\mu$ M and 0.0003  $\mu$ M.

#### Inhibition of TNF Production

The ability of the compounds or the pharmaceutically acceptable salts thereof to inhibit the production of TNF and, consequently, demonstrate their effectiveness for treating diseases involving the production of TNF is shown by the following in vitro assay:

Human mononuclear cells were isolated from anti-coagulated human blood using a one-step Ficoll-hypaque separation technique. The mononuclear cells were washed three times in Hanks balanced salt solution (HBSS) with divalent cations and resuspended to a density of  $2 \times 10^6$ /ml in HBSS containing 1% BSA. Differential counts determined using the Abbott Cell Dyn 3500 analyzer indicated that monocytes ranged from 17 to 24% of the total cells in these preparations.

180 $\mu$  of the cell suspension was aliquoted into flat bottom 96 well plates (Costar). Additions of compounds and LPS (100ng/ml final concentration) gave a final volume of 200 $\mu$ l. All conditions were performed in triplicate. After a four hour incubation at 37°C. in an humidified CO<sub>2</sub> incubator, plates were removed and centrifuged (10 minutes at approximately 250 x g) and the supernatants removed and assayed for TNF $\alpha$  using the R&D ELISA Kit.

For administration to humans for the inhibition of matrix metalloproteinases or the

- 32 -

production of tumor necrosis factor (TNF), a variety of conventional routes may be used including orally, parenterally and topically. In general, the active compound will be administered orally or parenterally at dosages between about 0.1 and 25 mg/kg body weight of the subject to be treated per day, preferably from about 0.3 to 5 mg/kg. However, some variation in dosage will necessarily occur depending on the condition of the subject being treated. The person responsible for administration will, in any event, determine the appropriate dose for the individual subject.

The compounds of the present invention can be administered in a wide variety of different dosage forms, in general, the therapeutically effective compounds of this invention are present in such dosage forms at concentration levels ranging from about 5.0% to about 70% by weight.

For oral administration, tablets containing various excipients such as microcrystalline cellulose, sodium citrate, calcium carbonate, dicalcium phosphate and glycine may be employed along with various disintegrants such as starch (and preferably corn, potato or tapioca starch), alginic acid and certain complex silicates, together with granulation binders like polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for tabletting purposes. Solid compositions of a similar type may also be employed as fillers in gelatin capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration, the active ingredient may be combined with various sweetening or flavoring agents, coloring matter or dyes, and, if so desired,

- 33 -

emulsifying and/or suspending agents as well, together with such diluents as water, ethanol, propylene glycol, glycerin and various like combinations thereof.

For parenteral administration (intramuscular, intraperitoneal, subcutaneous and intravenous use) a sterile injectable solution of the active ingredient is usually prepared. Solutions of a therapeutic compound of the present invention in either sesame or peanut oil or in aqueous propylene glycol may be employed. The aqueous solutions should be suitably adjusted and buffered, preferably at a pH of greater than 8, if necessary and the liquid diluent first rendered isotonic. These aqueous solutions are suitable intravenous injection purposes. The oily solutions are suitable for intraarticular, intramuscular and subcutaneous injection purposes. The preparation of all these solutions under sterile conditions is readily accomplished by standard pharmaceutical techniques well known to those skilled in the art.

The present invention is illustrated by the following examples, but it is not limited to the details thereof.

#### EXAMPLE 1

2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]-3-methylbutyramide

To a solution of D-valine benzyl ester hydrochloride (2.4 grams, 10 mmol) and triethylamine (2.5 grams, 3.5 mL, 25 mmol) in water (50 mL) and 1,4-dioxane (50 mL) is added 4-methoxybenzenesulfonyl chloride (2.3 grams, 11 mmol). The mixture was stirred at room temperature for 1 hour and then most of the solvent was removed by evaporation under vacuum. The mixture was diluted with ethyl acetate and was washed successively with dilute hydrochloric acid solution, water and brine. The organic solution was dried over magnesium sulfate and concentrated to leave N-(4-methoxybenzenesulfonyl)-D-valine benzyl ester as a

- 34 -

white solid, 3.6 grams (97%); m.p. 92-94°C.

N-(4-Methoxybenzenesulfonyl)-D-valine benzyl ester (1.50 grams, 4.0 mmol) was added to a suspension of sodium hydride (0.1 grams, 4.2 mmol) in dry dimethylformamide (20 mL) and, after 30 minutes, tert-butyl bromoacetate (0.8 mL, 4.2 mmol) was added. The resulting mixture was stirred overnight at room temperature and was then quenched by addition of saturated ammonium chloride solution (3 mL). The dimethylformamide was removed by evaporation under vacuum. The residue was taken up in ethyl acetate and washed with water and brine. After drying over magnesium sulfate, ethyl acetate was evaporated to leave an oil from which 2-(R)-2-[tert-butoxycarbonylmethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester, a clear oil (1.92 grams, 98%), was isolated using flash chromatography on silica gel eluting with 15% ethyl acetate in hexane.

To a cold (0°C.) solution of 2-(R)-2-[tert-butoxycarbonylmethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester (1.92 grams, 3.9 mmol) in methylene chloride (28 mL) was added trifluoroacetic acid (7 mL). The resulting solution was allowed to warm to room temperature and was stirred overnight. The methylene chloride and trifluoroacetic acid were evaporated under vacuum leaving 2-(R)-2-[carboxymethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester as a clear oil, 1.70 grams (100%).

To a solution of 2-(R)-2-[carboxymethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester (573 mg, 1.32 mmol) in methylene chloride (12 mL) were added sequentially triethylamine (0.46 mL, 3.28 mmol), morpholine (0.127 mL, 1.46 mmol) and (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (646 mg, 1.46 mmol). The mixture was stirred at room temperature overnight and then

- 35 -

diluted with ethyl acetate. The solution was washed with 0.5 N hydrochloric acid solution and brine, dried over magnesium sulfate and concentrated under vacuum. The residue was chromatographed on silica gel using 40% ethyl acetate in hexane affording 2-(R)-2-[(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-oxoethyl)amino]-3-methylbutyric acid benzyl ester as a clear oil, 590 mg (89%).

To a solution of 2-(R)-2-[(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]-3-methylbutyric acid benzyl ester (590 mg, 1.17 mmol) in ethanol (50 mL) was added 10% palladium on activated carbon (200 mg). The mixture was agitated under 3 atmospheres hydrogen in a Parr shaker for 2 hours. The catalyst was removed by filtration through nylon (pore size 0.45  $\mu$ m) and the solvent was evaporated leaving 2-(R)-2-[(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]-3-methylbutyric acid as a white foam, 485 mg (100%).

To a solution of 2-(R)-2-[(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]-3-methylbutyric acid (485 mg, 1.17 mmol) in methylene chloride (12 mL) were added sequentially triethylamine (0.52 mL, 3.71 mmol), O-benzylhydroxylamine hydrochloride (205 mg, 1.28 mmol) and (benzotriazol-1-yl)oxy)tris(dimethylamino)phosphonium hexafluorophosphate (570 mg, 1.29 mmol). The mixture was stirred at room temperature overnight and then diluted with ethyl acetate. The solution was washed sequentially with 0.5 N hydrochloric acid solution, water, saturated sodium hydrogen carbonate solution and brine, dried over magnesium sulfate and concentrated under vacuum. The residue was chromatographed on silica gel using 20% hexane in ethyl acetate to afford 2-(R)-N-benzyloxy-2-[(4-methoxybenzenesulfonyl)(2-

- 36 -

morpholin-4-yl-2-oxoethyl)amino]-3-methylbutyramide as a white foam, 510 mg (84%).

To a solution of 2-(R)-N-benzyloxy-2-[(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]-3-methylbutyramide (510 mg, 0.98 mmol) in methanol (50 mL) was added 5% palladium on activated carbon (120 mg). The mixture was agitated under 2 atmospheres hydrogen in a Parr shaker for 2 hours. The catalyst was removed by filtration through nylon (pore size 0.45  $\mu$ m) and the solvent was evaporated leaving 2-(R)-N-hydroxy-2-[(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]-3-methylbutyramide as a white solid, 418 mg (99%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  10.3 (br s, 1H), 7.90 (br s, 1H, overlapped), 7.86 (d, J = 8.8 Hz, 2H, overlapped), 6.94 (d, J = 8.8 Hz, 2H), 4.39 (d, J = 17.1 Hz, 1H), 4.09 (d, J = 17.1, 1H), 3.84 (s, 3H), 3.80-3.48 (m, 9H), 2.20-1.95 (m, 1H), 0.82 (d, J = 6.5 Hz, 3H), 0.45 (d, J = 6.5 Hz, 3H); MS (LSIMS): m/z 430 (M+H).

20

#### EXAMPLE 2

##### 2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)(3-morpholin-4-yl-3-oxopropyl)amino]-3-methylbutyramide

To a solution of N-(4-methoxybenzenesulfonyl)-D-valine benzyl ester (2.2 grams, 5.83 mmol) in dry dimethylformamide (40 mL) were added cesium carbonate (2.3 grams, 7.1 mmol) and 1-iodo-3-butene (1.3 grams, 7.1 mmol). The mixture was stirred at room temperature overnight and was then poured into water. The mixture was extracted twice with ether and the combined ether extracts were washed with brine, dried over magnesium sulfate and concentrated under reduced pressure. The residue was taken up in 20% ethyl acetate/hexane; starting material N-(4-methoxybenzenesulfonyl)-D-valine benzyl ester (1.5 g) crystallized from the mixture and was recovered by filtration. The filtrate was concentrated under vacuum and the residue was chromatographed on silica

- 37 -

gel using 20% ethyl acetate/hexane as eluent to provide 2-(R)-2-[but-3-enyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester as a clear oil, 404 mg (16%).

5 To a mixture of 2-(R)-2-[but-3-enyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester (780 mg, 1.81 mmol) and ruthenium (III) chloride hydrate (10 mg, 0.048 mmol) in acetonitrile (6 mL), carbon tetrachloride (6 mL) and water (8 mL) was  
10 added sodium periodate (1.7 grams, 7.9 mmol). After stirring at room temperature for 2 hours, the mixture was diluted with methylene chloride and filtered through diatomaceous earth. The organic layer was  
15 separated, washed with dilute hydrochloric acid solution and brine, dried over magnesium sulfate and concentrated to leave 2-(R)-2-[2-carboxyethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester as an oil, 710 mg (87%).

Alternatively, the intermediate 2-(R)-2-[2-carboxyethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester was prepared by the  
20 following higher yielding procedure:

N-(4-Methoxybenzenesulfonyl)-D-valine benzyl ester (18.8 grams, 49.8 mmol) was added to a suspension  
25 of sodium hydride (1.3 grams, 54 mmol) in dry dimethylformamide (200 mL) and, after 1.5 hours, a solution of allyl bromide (4.7 mL, 54 mmol) was added. The resulting mixture was stirred overnight at room temperature and was then quenched by addition of  
30 saturated ammonium chloride solution. The dimethylformamide was removed by evaporation under vacuum. The residue was taken up in ether and washed with water and brine. After drying over magnesium sulfate, ether was evaporated to leave an oil from  
35 which 2-(R)-2-[(4-methoxybenzenesulfonyl)prop-2-enylamino]-3-methylbutyric acid benzyl ester, a clear oil (18.1 grams, 87%), was isolated using flash

- 38 -

chromatography on silica gel eluting with 10% ethyl acetate in hexane and then 20% ethyl acetate in hexane.

To a 1 M solution of borane/disulfide complex in methylene chloride (1.45 mL, 2.9 mmol) was added a solution of 2-(R)-2-[(4-methoxybenzenesulfonyl)prop-2-enylamino]-3-methylbutyric acid benzyl ester (3.6 grams, 8.6 mmol) in methylene chloride (8 mL). The solution was stirred at room temperature for 4 hours at which time more 1 M solution of borane/disulfide complex in methylene chloride (2.0 mL, 4.0 mmol) was added. The mixture was stirred at room temperature overnight and was then added dropwise to a mechanically stirred solution of chromium trioxide (5.1 grams, 51.6 mole) in acetic acid (31 mL) and water (3.5 mL) while keeping the internal temperature between -5°C. and 10°C. After stirring at room temperature overnight, the mixture was diluted with water and extracted with methylene chloride. The extract was washed with brine, dried (magnesium sulfate) and concentrated. The residue was chromatographed on silica gel eluting successively with chloroform and 2% methanol in chloroform to afford 2-(R)-2-[2-carboxyethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl as an oil (2.42 grams, 63%).

To a solution of 2-(R)-2-[2-carboxyethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester (710 mg, 1.58 mmol) in methylene chloride (15 mL) were added sequentially triethylamine (0.47 mL, 3.35 mmol), morpholine (0.15 mL, 1.72 mmol) and (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (769 mg, 1.74 mmol). The mixture was stirred at room temperature overnight and then diluted with methylene chloride. The solution was washed with 0.5 N hydrochloric acid solution and brine, dried over magnesium sulfate and concentrated under vacuum. The solid residue was chromatographed on silica gel using 20% hexane in ethyl acetate affording



- 39 -

2-(R)-2-[(4-methoxybenzenesulfonyl)(3-morpholin-4-yl-3-oxopropyl)amino]-3-methylbutyric acid benzyl ester as a clear oil, 725 mg (88%).

To a solution of 2-(R)-2-[(4-methoxybenzenesulfonyl)(3-morpholin-4-yl-3-oxopropyl)amino]-3-methylbutyric acid benzyl ester (725 mg, 1.40 mmol) in ethanol (35 mL) was added 10% palladium on activated carbon (50 mg). The mixture was agitated under 3 atmospheres hydrogen in a Parr shaker for 3 hours. The catalyst was removed by filtration through nylon (pore size 0.45  $\mu$ m) and the solvent was evaporated leaving 2-(R)-2-[(4-methoxybenzenesulfonyl)(3-morpholin-4-yl-3-oxopropyl)amino]-3-methyl-butylric acid as a white solid, 540 mg (90%).

To a solution of 2-(R)-2-[(4-methoxybenzenesulfonyl)(3-morpholin-4-yl-3-oxopropyl)amino]-3-methylbutyric acid (540 mg, 1.26 mmol) and 1-hydroxybenztriazole hydrate (205 mg, 1.33 mmol) in dry dimethylformamide (12 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (289 mg, 1.51 mmol). After stirring for 30 minutes, hydroxylamine hydrochloride (350 mg, 5.04 mmol) and then triethylamine (1.0 mL, 7.17 mmol) were added. The mixture was stirred at room temperature overnight and then diluted with ethyl acetate. The solution was washed sequentially with water, 0.5 N hydrochloric acid solution and brine. The solution was then dried over magnesium sulfate and concentrated under vacuum to leave a white foam. The material was dissolved in toluene, filtered and concentrated. The residue was triturated with ether to afford 2-(R)-N-hydroxy-2[(4-methoxybenzenesulfonyl)(3-morpholin-4-yl-3-oxopropyl)amino]-3-methylbutyramide as a solid, 200 mg (36%);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  9.35 (br s, 1H), 7.73 (d, J = 8.9 Hz, 2H), 6.95 (d, J = 8.9 Hz, 2H), 3.86 (s, 3H), 3.83-3.73 (m, 1H), 3.70-3.52 (m, 7H), 3.46-3.43 (m,

- 40 -

2H), 3.41-3.29 (m, 1H), 2.92-2.69 (m, 2H), 2.30-2.17 (m, 1H), 0.84 (d, J = 6.5 Hz, 3H), 0.41 (d, J = 6.5 Hz, 3H); MS (particle beam): m/z 444 (M+H), 428, 383, 329; HRMS calculated for C<sub>19</sub>H<sub>30</sub>N<sub>3</sub>O<sub>7</sub>S (M+H): 444.1804,

5 Found: 464.1818.

The title compounds of Examples 3-6 were prepared by a method analogous to that described in Example 2 using 2-(R)-2-[2-carboxyethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid  
10 benzyl ester as the starting material which is coupled with the amine indicated.

#### EXAMPLE 3

2-(R)-2-[(2-Benzylcarbamoyl)ethyl](4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide  
15

Coupled with benzylamine; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>):  
δ 10.72 (s, 1H), 8.89 (s, 1H), 8.39 (m, 1H), 7.74 (d, J = 9.0 Hz, 2H), 7.32-7.21 (m, 5H), 7.05 (d, J = 9.0 Hz, 2H), 4.21 (d, J = 5.9 Hz, 2H), 4.02-3.87 (m, 1H), 3.82  
20 (s, 3H), 3.63 (d, J = 10.8 Hz, 1H), 3.29-3.17 (m, 1H), 2.71-2.57 (m, 1H), 2.52-2.40 (m, 1H), 2.06-1.94 (m, 1H), 0.77 (d, J = 6.6 Hz, 3H), 0.74 (d, J = 6.5 Hz, 3H); MS (LSIMS): m/z 464 (M+H); HRMS calculated for C<sub>22</sub>H<sub>30</sub>N<sub>3</sub>O<sub>6</sub>S (M+H): 464.1855. Found: 464.1832.

25

#### EXAMPLE 4

2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)(2-[(pyridin-3-ylmethyl)carbamoyl]ethyl)amino]-3-methylbutyramide

Coupled with 3-pyridylmethylamine: <sup>1</sup>H NMR (DMSO-d<sub>6</sub>):  
30 δ 10.72 (s, 1H), 8.89 (s, 1H), 8.49-8.42 (m, 3H), 7.73 (d, J = 8.9 Hz, 2H), 7.63-7.60 (m, 1H), 7.32 (dd, J = 4.8, 7.8 Hz, 1H), 7.05 (d, J = 8.9 Hz, 2H), 4.23 (d, J = 5.8 Hz, 2H), 4.00-3.88 (m, 1H), 3.81 (s, 3H), 3.62 (d, J = 10.8 Hz, 1H), 3.27-3.17 (m, 1H),  
35 2.69-2.58 (m, 1H), 2.52-2.41 (m, 1H), 2.07-1.94 (m, 1H), 0.76 (d, J = 6.5 Hz, 3H), 0.72 (d, J = 6.4 Hz, 3H); MS (LSIMS): m/z 465 (M+H).

- 41 -

EXAMPLE 52-(R)-N-Hydroxy-2-([4-methoxybenzenesulfonyl]  
[2-(methylpyridin-3-ylmethylcarbamoyl)ethyl]amino)-3-  
methylbutyramide

5 Coupled with 3-(N-methylaminomethyl)pyridine:  
<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 10.75 (br s, 1H), 8.92 (s, 1H),  
 8.52-8.29 (m, 2H), 7.75 (d, J = 8.8 Hz, 1.4 H), 7.67  
 (d, J = 8.8 Hz, 0.6 H), 7.62-7.58 (m, 1H), 7.42-7.32  
 (m, 1H), 7.06 (d, J = 8.8 Hz, 1.4 H), 7.01 (d, J = 8.8  
 10 Hz, 0.6 H), 4.55-4.41 (m, 2H), 3.94-3.82 (m, 1H), 3.81  
 (s, 2.1 H), 3.80 (s, 0.9 H), 3.68-3.60 (m, 1H), 3.33-  
 3.19 (m, 1H), 2.90-2.50 (m, 2H), 2.88 (s, 2.1 H  
 overlapped), 2.79 (s, 0.9 H), 2.05-1.80 (m, 1H),  
 0.79-0.63 (m, 6H): MS (thermospray): m/z 479  
 15 (M+H), 364.

EXAMPLE 64-(3-[(1-(R)-1-Hydroxycarbamoyl-2-  
methylpropyl)(4-methoxybenzenesulfonyl)amino]propionyl)  
piperazine-1-carboxylic acid, tert-butyl ester

20 Coupled with tert-butyl-1-  
 piperazinecarboxylate: <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 10.77 (br  
 s, 1H), 8.93 (s, 1H), 7.74 (d, J = 8.9 Hz, 2H), 7.06  
 (d, J = 8.9 Hz, 2H), 3.90-3.80 (m, 1H), 3.82 (s, 3H,  
 overlapped), 3.64 (d, J = 10.8 Hz, 1H), 3.60-3.16 (m,  
 25 9H), 2.80-2.71 (m, 1H), 2.59-2.47 (m, 1H), 2.03-1.91  
 (m, 1H), 1.39 (s, 9H), 0.77 (d, J = 6.5 Hz, 3H), 0.71  
 (d, J = 6.5, 3H); MS (thermospray): m/z 543 (M+H),  
 443, 382, 328.

EXAMPLE 7

30 2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)  
 (3-oxo-3-piperazin-1-ylpropyl)amino]-3-methylbutyramide  
 hydrochloride

A solution of 4-(3-[(1-(R)-1-  
 hydroxycarbamoyl-2-methylpropyl)(4-  
 35 methoxybenzenesulfonyl)amino]propionyl)piperazine-1-  
 carboxylic acid, tert-butylester [Example 6] (430 mg,  
 0.79 mmol) in methylene chloride (11 mL) was cooled to

- 42 -

0°C. Hydrogen chloride gas was then bubbled through the solution for about 0.5 minute. The solution was allowed to warm to room temperature with stirring over 1 hour. Volatiles were removed by evaporation and the residue was filtered washing with methylene chloride to collect solid 2-(R)-N-hydroxy-2-[(4-methoxybenzenesulfonyl)(3-oxo-3-piperazin-1-ylpropyl)amino]-3-methylbutyramide hydrochloride, 375 mg (99%). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 10.78 (br s, 1H), 9.16 (br s, 1H), 7.74 (d, J = 8.8 Hz, 2H), 7.07 (d, J = 8.9 Hz, 2H), 3.82 (s, 3H), 3.62 (br s, 4H), 3.38-3.18 (m, 1H), 3.16-3.07 (br s, 2H), 3.07-2.98 (br s, 2H), 2.83-2.73 (m, 1H), 2.65-2.53 (m, 1H), 2.06-1.90 (m, 1H), 0.76 (d, J = 6.5 Hz, 3H), 0.72 (d, J = 6.5 Hz, 3H). A broad water peak between δ 4.0 and 3.5 obscured some signals from this compound; MS (thermospray): m/z 443 (M+H), 382, 328.

#### EXAMPLE 8

2-(R)-2-[(Benzylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide

To a solution of 2-(R)-2-[carboxymethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester (example 1) (905 mg, 2.08 mmol) in methylene chloride (18 mL) were added sequentially triethylamine (0.72 mL, 5.14 mmol), benzylamine (0.25 mL, 2.29 mmol) and (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (1.01 grams, 2.28 mmol). The mixture was stirred at room temperature overnight and then diluted with ethyl acetate. The solution was washed with 0.5 N hydrochloric acid solution and brine, dried over magnesium sulfate and concentrated under vacuum. The residue was chromatographed on silica gel using a 2:5:16 ratio of methylene chloride/ethyl acetate/hexane affording 2-(R)-2-[(benzylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid

- 43 -

benzyl ester as a clear oil, 933 mg (86%).

To a solution of 2-(R)-2-

[(benzylcarbamoylmethyl)(4-methoxybenzenesulfonyl)-  
amino]-3-methylbutyric acid benzyl ester (933 mg, 1.17  
5 mmol) in ethanol (50 mL) was added 10% palladium on  
activated carbon (85 mg). The mixture was agitated  
under 3 atmospheres hydrogen in a Parr shaker for 4  
hours. The catalyst was removed by filtration through  
nylon (pore size 0.45  $\mu$ m) and the solvent was  
10 evaporated leaving 2-(R)-2-[(benzylcarbamoylmethyl)(4-  
methoxybenzenesulfonyl)amino]-3-methylbutyric acid as a  
white foam, 755 mg (98%).

To a solution of 2-(R)-2-

[(benzylcarbamoylmethyl)(4-  
15 methoxybenzenesulfonyl)amino]-3-methylbutyric acid (655  
mg, 1.51 mmol) and 1-hydroxybenztriazole hydrate (224  
mg, 1.46 mmol) in dry dimethylformamide (15 mL) was  
added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide  
hydrochloride (316 mg, 1.65 mmol). After stirring for  
20 30 minutes, hydroxylamine hydrochloride (416 mg, 6.0  
mmol) and then N-methylmorpholine (0.99 mL, 9.0 mmol)  
were added. The mixture was stirred at room  
temperature overnight and then diluted with ethyl  
acetate. The solution was washed sequentially with  
25 water, 0.5 N hydrochloric acid solution and brine. The  
solution was then dried over magnesium sulfate and  
concentrated under vacuum to leave a white foam which  
was chromatographed on silica gel eluting with ethyl  
acetate to afford 2-(R)-2-[(benzylcarbamoylmethyl)(4-  
30 methoxybenzene-sulfonyl)amino]-N-hydroxy-3-  
methylbutyramide as a white foam, 570 mg (84%);  $^1\text{H}$  NMR  
(DMSO- $d_6$ ):  $\delta$  10.75 (br s, 1H), 8.90 (s, 1H), 8.47 (m,  
1H), 7.85 (d, J = 8.9 Hz, 2H), 7.83-7.19 (m, 5H), 7.04  
(d, J = 8.9 Hz, 2H), 4.37 (d, J = 11.4 Hz, 1H), 4.28  
35 (d, J = 5.9 Hz, 2H), 3.89 (d, J = 11.4 Hz, 1H), 3.82  
(s, 3H), 3.45 (d, J = 10.3 Hz, 1H), 1.90-1.79 (m, 1H),  
0.73 (d, J = 6.3 Hz, 6H); MS (LSIMS): m/z 450 (M+H).

- 44 -

EXAMPLE 92-(R)-2-[(Benzylmethylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide

5 To a solution of 2-(R)-2-[carboxymethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester (Example 1) (1.05 grams, 2.41 mmol) in methylene chloride (20 mL) were added sequentially triethylamine (0.84 mL, 6.0 mmol), N-benzylmethylamine  
10 (0.34 mL, 2.63 mmol) and (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (1.17 grams, 2.69 mmol). The mixture was stirred at room temperature overnight and then diluted with ethyl acetate. The solution was  
15 washed with 0.5 N hydrochloric acid solution and brine, dried over magnesium sulfate and concentrated under vacuum. The residue was chromatographed on silica gel using 35% ethyl acetate in hexane (plus a small amount of methylene chloride to load the sample on the column)  
20 affording 2-(R)-2-[benzylmethylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester as a clear oil, 1.14 grams (88%).

To a solution of 2-(R)-2-[(benzylmethylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid  
25 benzyl ester (1.14 grams, 2.12 mmol) in ethanol (100 mL) was added 10% palladium on activated carbon (400 mg). The mixture was agitated under 3 atmospheres hydrogen in a Parr shaker for 3 hours. The catalyst  
30 was removed by filtration through nylon (pore size 0.45  $\mu$ m) and the solvent was evaporated leaving 2-(R)-2-[(benzylmethylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid as a white foam, 902 mg (95%).

35 To a solution of 2-(R)-2-[(benzylmethylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid (902

- 45 -

mg, 2.01 mmol) in methylene chloride (20 mL) were added sequentially triethylamine (0.90 mL, 6.42 mmol), O-allylhydroxylamine hydrochloride (242 mg, 2.21 mmol) and (benzotriazol-1-yloxy)tris-

5 (dimethylamino)phosphonium hexafluorophosphate (978 mg, 2.21 mmol). The mixture was stirred at room temperature overnight and then diluted with ethyl acetate. The solution was washed with 0.5 N hydrochloric acid solution and brine, dried over

10 magnesium sulfate and concentrated under vacuum. The residue was chromatographed on silica gel using 40% hexane in ethyl acetate to afford 2-(R)-N-allyloxy-2-[(benzylmethylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyramide as an oil,

15 1.008 grams (100%).

To a solution of 2-(R)-N-allyloxy-2-[(benzylmethyl-carbamoylmethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyramide (500 mg, 0.99 mmol) in methylene chloride (40 mL) was added

20 bis(triphenylphosphine)palladium (II) chloride (280 mg, 0.4 mmol) and then, dropwise, tributyltinhydride (0.43 mL, 2.2 mmol). The solution was stirred at room temperature for 1 hour, diluted with methylene chloride, washed with 1 N hydrochloric acid solution,

25 dried over magnesium sulfate and concentrated. The residue was taken up in ethyl acetate and filtered to remove a solid. After concentration, the filtrate was chromatographed on silica gel eluting with chloroform and then 2% methanol in chloroform to afford 2-(R)-2-

30 [(benzylmethylcarbamoylmethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide as a white solid (340 mg, 74%). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 10.66 (br s, 1H), 8.87 (br s, 0.6 H), 8.84 (s, 0.4 H), 7.91 (d, J = 8.9 Hz, 1.2 H), 7.78 (d, J = 8.9 Hz, 0.8 H), 7.43-7.21

35 (m, 5H), 7.05 (d, J = 8.9 Hz, 1.2 H), 7.00 (d, J = 8.9 Hz, 0.8 H) 4.72 (d, J = 17.7 Hz, 0.4 H), 4.70 (d, J = 17.7 Hz, 0.6 H), 4.59-4.42 (m, 1H), 4.25 (d, J = 17.8

- 46 -

Hz, 0.6H), 4.07 (d, J = 17.7 Hz, 0.4H), 3.82 (s, 3H), 3.46-3.40 (m, 1H), 2.91 (s, 1.8H), 2.83 (s, 1.2 H), 1.92-1.70 (m, 1H), 0.75-0.69 (m, 6H); MS (thermospray): m/z 464 (M+H), 307, 239.

5           The title compounds of Examples 10-11 were prepared by a method analogous to that described in Example 9 using 2-(R)-2-[carboxymethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester (Example 1) as the starting material which  
10 is coupled with the amine indicated.

EXAMPLE 10

2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)-(2-morpholin-4-ylethylcarbamoyl)methyl]amino)-3-methylbutyramide

15           Coupled with 4-(2-aminoethyl)morpholine: <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): δ 10.74 (br s, 1H), 8.90 (br s, 1H), 7.84 (br s, 1H, overlapped), 7.84 (d, J = 8.8 Hz, 2H), 7.06 (d, J = 8.8 Hz, 2H), 4.33 (d, J = 17.5 Hz, 1H), 3.83 (s, 3H), 3.78 (d, J = 17.5 Hz, 1H), 3.57-3.54  
20 (m, 4H), 3.49 (d, J = 10.2 Hz, 1H), 3.28-3.06 (m, 2H), 2.34-2.30 (m, 6H), 1.93-1.77 (m, 1H), 0.77-0.74 (m, 6H).

EXAMPLE 11

2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)-(2-oxo-2-pyrrolidin-1-ylethyl)amino]-3-methylbutyramide

25           Coupled with pyrrolidine: <sup>1</sup>H NMR (CD<sub>3</sub>OD): δ 7.90 (d, J = 8.9 Hz, 2H), 7.04 (d, J = 8.9 Hz, 2H), 4.50 (d, J = 17.6 Hz, 1H), 4.15 (d, J = 17.6 Hz, 1H),  
30 3.87 (s, 3H), 3.56-3.39 (m, 5H), 2.07-1.82 (m, 5H), 0.83 (d, J = 6.6 Hz, 3H), 0.73 (d, J = 6.6 Hz, 3H); MS (thermospray): m/z 414 (M+1); HRMS calculated for C<sub>18</sub>H<sub>28</sub>N<sub>3</sub>O<sub>6</sub>S (M+H): 414.1699. Found 414.1703.



- 47 -

EXAMPLE 122-[Dimethylcarbamoylmethyl(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide

5           A solution of 2-(R)-2-[carboxymethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester (Example 1) (1.89 grams, 4.34 mmol) in thionyl chloride (25 mL) was heated at reflux for 1 hour. After cooling, the excess thionyl chloride was  
10 evaporated. The residue was taken up in methylene chloride (50 mL) and the solution was cooled in an ice bath. Dimethylamine gas was slowly bubbled through the solution for 1 hour. After evaporation of the solvent, the residue was taken up in ethyl acetate, washed with  
15 1 N hydrochloric acid solution, water and brine, dried over magnesium sulfate and concentrated to leave dimethylcarbamoylmethyl(4-methoxybenzenesulfonyl)amino-3-methylbutyric acid benzyl ester as an oil, 1.77 grams (88%).

20           To a solution of dimethylcarbamoylmethyl(4-methoxybenzenesulfonyl)amino-3-methylbutyric acid benzyl ester (1.77 grams, 3.83 mmol) in ethanol (100 mL) was added 10% palladium on activated carbon (644 mg). The mixture was agitated under 3 atmospheres  
25 hydrogen in a Parr shaker for 1.5 hours. The catalyst was removed by filtration through nylon (pore size 0.45  $\mu$ m) and the solvent was evaporated leaving dimethylcarbamoylmethyl(4-methoxybenzenesulfonyl)amino-3-methylbutyric acid as a white foam, 1.42 grams (100%).

30           To a solution of dimethylcarbamoylmethyl(4-methoxybenzenesulfonyl)amino-3-methylbutyric acid (1.42 grams, 3.81 mmol) and 1-hydroxybenztriazole hydrate (687 mg, 4.48 mmol) in dry dimethylformamide (7 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide  
35 hydrochloride (974 mg, 5.08 mmol). After stirring for 30 minutes, hydroxylamine hydrochloride (1.17 grams, 16.8 mmol) and then N-methylmorpholine (2.8 mL, 25.5

- 48 -

mmol) were added. The mixture was stirred at room temperature overnight and then concentrated under vacuum. The residue was taken up in ethyl acetate and the resulting solution was washed sequentially with water, 0.5 N hydrochloric acid solution and brine. The solution was then dried over magnesium sulfate and concentrated under vacuum to leave an oil which was chromatographed on silica gel eluting successively with ethyl acetate, 5% methanol in chloroform and 10% methanol in chloroform to afford 2-[dimethyl-carbamoylmethyl(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide as a white solid, 390 mg (26%).  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  10.70 (br s, 1H), 8.89 (s, 1H), 7.80 (d,  $J = 8.9$  Hz, 2H), 7.10 (d,  $J = 8.9$  Hz, 2H), 4.62 (d,  $J = 17.7$  Hz, 1H), 4.14 (d,  $J = 17.7$  Hz, 1H), 3.84 (s, 3H), 3.40 (d,  $J = 10.4$  Hz, 1H), 2.97 (s, 3H), 2.82 (s, 3H), 1.88-1.72 (m, 1H), 0.72 (d,  $J = 6.5$  Hz, 6H); MS (thermospray):  $m/z$  388 ( $M+1$ ); HRMS calculated for  $\text{C}_{16}\text{H}_{26}\text{N}_3\text{O}_6\text{S}$  ( $M+H$ ): 388.1542 Found: 388.1592.

### EXAMPLE 13

2-(R)-2-N-Hydroxy-2-((4-methoxybenzenesulfonyl)((pyridin-3-ylmethyl)carbamoyl)methyl)amino)-3-methylbutyramide

25                    2-(R)-N-Hydroxy-2-((4-methoxybenzenesulfonyl)((pyridin-3-ylmethyl)carbamoyl)methyl)amino)-3-methylbutyramide was prepared by a procedure similar to that of Example 12 starting with 2-(R)-2-[carboxymethyl(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester (Example 1) and coupling this to 3-pyridylmethylamine via the acid chloride.  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  8.55-8.53 (m, 1H), 8.43-8.40 (m, 1H), 7.90-7.82 (m, 1H, overlapped), 7.86 (d,  $J = 8.9$  Hz, 2H), 7.40 (dd,  $J = 4.8, 7.8$  Hz, 1H), 7.04 (d,  $J = 8.9$  Hz, 2H), 4.50 (d,  $J = 17.5$  Hz, 1H), 4.39 (d,  $J = 17.5$  Hz, 1H), 4.32 (d,  $J = 17.7$  Hz, 1H), 4.02 (d,  $J = 17.7$  Hz,

- 49 -

1H), 3.87 (s, 3H), 3.60 (d, J = 10.3 Hz, 1H), 2.08-1.93 (m, 1H), 0.85 (d, J = 6.5 Hz, 3H), 0.70 (d, J = 6.5 Hz, 3H); MS (thermospray): m/z 451 (M+H), 336, 320.

EXAMPLE 14

5                    N-Hydroxy-[(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]acetamide

To a solution of iminoacetic acid disodium salt monohydrate (5.0 grams, 25.6 mmol) in dioxane (50 ml) and water (50 ml) was added triethylamine (5.3 ml, 38 mmol) followed by 4-methoxybenzenesulfonyl chloride (5.8 grams, 28.0 mmol). The mixture was stirred overnight at room temperature and diluted with methylene chloride. The solution was washed with 1 N hydrochloric acid solution, water and brine, dried over  
10                    magnesium sulfate and concentrated under vacuum leaving  
15                    [carboxymethyl(4-methoxybenzenesulfonyl)amino]acetic acid as a white solid, 3.83 grams (49%).

[Carboxymethyl(4-methoxybenzenesulfonyl)amino]acetic acid (0.5 grams, 1.65 mmol) in acetic anhydride (15 mL) was dissolved in acetic anhydride by gentle warming. The resulting solution was stirred at room temperature overnight. The acetic anhydride was removed by evaporation under vacuum; the residue was dissolved in methylene chloride  
20                    and morpholine (0.16 mL, 1.82 mmol) was added. The mixture was stirred overnight at room temperature and then concentrated under vacuum. The residue was dissolved in ethyl acetate, washed with 1 N hydrochloric acid solution, water and brine, dried over  
25                    magnesium sulfate and concentrated to afford [(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]acetic acid as an oil, 0.33 grams (54%).

To a solution of [(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)-  
35                    amino]acetic acid (0.33 grams, 0.89 mmol) in methylene chloride (10 mL) were added sequentially triethylamine (0.43 mL, 3.1 mmol), O-benzylhydroxylamine

- 50 -

hydrochloride (0.15 grams, 0.94 mmol) and (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (0.43 grams, 0.97 mmol). The mixture was stirred at room temperature overnight and then diluted with ethyl acetate. The solution was washed sequentially with 0.5 N hydrochloric acid solution, water and brine, dried over magnesium sulfate and concentrated under vacuum. The residue was chromatographed on silica gel using ethyl acetate to afford N-benzyloxy-[(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]acetamide as a white solid, 0.33 grams (78%).

To a solution of N-benzyloxy-[(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]acetamide (0.33 grams, 0.69 mmol) in methanol (35 mL) was added 5% palladium on activated carbon (85 mg). The mixture was agitated under 2 atmospheres hydrogen in a Parr shaker for 1.5 hours. The catalyst was removed by filtration through nylon (pore size 0.45  $\mu$ m) and the solvent was evaporated. The residue was chromatographed on silica gel eluting with 5% methanol in methylene chloride to afford N-methoxy-[(4-methoxybenzenesulfonyl)(2-morpholin-4-yl-2-oxoethyl)amino]acetamide as a white solid, 65 mg (24%);  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ ):  $\delta$  7.82 (d, J = 9.0 Hz, 2H), 7.08 (d, J = 9.0 Hz, 2H), 4.24 (s, 2H), 3.88 (s, 3H), 3.84 (s, 2H), 3.68-3.64 (m, 4H), 3.58-3.50 (m, 4H); MS (thermospray): m/z 388 (M+1), 387 (M); HRMS calculated for  $\text{C}_{16}\text{H}_{22}\text{N}_3\text{O}_7\text{S}$  (M+H): 388.1178, Found 338.1180.

The title compounds of Examples 15-16 were prepared by a method analogous to that described in Example 14 using [carboxymethyl(4-methoxybenzenesulfonyl)amino]acetic acid as the starting material which, after treatment with acetic anhydride, is coupled with the amine indicated.

- 51 -

EXAMPLE 15N-Hydroxy-[(4-methoxybenzenesulfonyl)(2-oxo-2-pyrrolidin-1-ylethyl)amino]acetamide

Coupled with pyrrolidine:  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  
5  $\delta$  11.26 (br s, 1H), 8.89 (s, 1H), 7.81 (d,  $J = 8.9$  Hz, 2H), 7.10 (d,  $J = 8.9$  Hz, 2H), 4.09 (s, 2H), 3.85 (s, 3H), 3.74 (s, 2H), 3.45-3.25 (m, 4H), 1.93-1.72 (m, 4H); MS (thermospray):  $m/z$  372 ( $M+1$ ): Analysis calculated for  $\text{C}_{15}\text{H}_{21}\text{N}_3\text{O}_6\text{S}$ : C, 48.51; H, 5.70; N, 11.31. Found: C, 48.51; H, 5.82; N, 11.24.

EXAMPLE 162-[Dimethylcarbamoylmethyl(4-methoxybenzenesulfonyl)amino]-N-hydroxyacetamide

Coupled with dimethylamine: mp:  $170^\circ\text{C}$ .  
15 (dec.);  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  10.69 (br s, 1H), 8.88 (s, 1H), 7.91 (d,  $J = 8.9$  Hz, 2H), 7.06 (d,  $J = 8.9$  Hz, 2H), 4.19 (s, 2H), 3.85 (s, 3H), 3.73 (s, 2H), 2.94 (s, 3H), 2.84 (s, 3H); MS (thermospray):  $m/z$  346 ( $M+1$ ); Analysis calculated for  $\text{C}_{13}\text{H}_{19}\text{N}_3\text{O}_6\text{S}$ : C, 45.21; H, 5.55; N, 12.17. Found: C, 44.93; H, 5.61; N, 12.03.

EXAMPLE 172-(R)-2-[(2-Carbamoylethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide

25 To a solution of 2-(R)-2-[(2-carboxylethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester (example 2) (900 mg., 2.0 mmol) in methylene chloride (10 mL) was added thionyl chloride (0.16 mL, 2.2 mmol). The reaction mixture was stirred  
30 for 1.5 hours at room temperature and then concentrated in vacuo. After dissolving the residue in methylene chloride (10 mL), ammonia gas was bubbled through the solution for 0.5 minutes. The solution was stirred at room temperature overnight and was concentrated under  
35 vacuum. Flash chromatography of the residue on silica gel eluting with 2% methanol in chloroform provided 2-(R)-2-[(2-carbamoylethyl)(4-

- 52 -

methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyric acid benzyl ester as a clear oil (275 mg, 31%).

To a solution of 2-(R)-2-[(2-carbamoylethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyric acid benzyl ester (275 mg, 0.61 mmol) in ethanol (15 mL) was added 10% palladium on activated carbon (30 mg). The mixture was agitated under 3 atmospheres hydrogen in a Parr shaker for 5 hours. The catalyst was removed by filtration through diatomaceous earth and the solvent was evaporated leaving 2-(R)-2-[(2-carbamoylethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyric acid as a white foam, 211 mg (96%).

To a solution of 2-(R)-2-[(2-carbamoylethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyric acid (205 mg, 0.57 mmol) and 1-hydroxybenzotriazole hydrate (85 mg, 0.55 mmol) in dry dimethylformamide (5 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (120 mg, 0.63 mmol). After stirring for 30 minutes, hydroxylamine hydrochloride (158 mg, 2.3 mmol) and then N-methylmorpholine (0.37 mL, 3.4 mmol) were added. The mixture was stirred at room temperature overnight and then diluted with ethyl acetate. The solution was washed with water and brine. The solution was then dried over magnesium sulfate and concentrated under vacuum to leave an oil which was chromatographed on silica gel eluting with 2% methanol in chloroform to afford 2-(R)-2-[(2-carbamoylethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide as a white solid, 45 mg (21%);  $^1\text{H}$  NMR (DMSO- $d_6$ ):  $\delta$  10.74 (br s, 1H), 8.91 (br s, 1H), 7.74 (d, J = 8.8 Hz, 2H), 7.33 (br s, 1H), 7.07 (d, J = 8.8 Hz, 2H), 6.79 (br s, 1H), 3.93-3.82 (m, 1H, overlapped), 3.83 (s, 3H), 3.64 (d, J = 10.7 Hz, 1H), 3.25-3.12 (m, 1H), 2.62-2.48 (m, 1H), 2.42-2.30 (m, 1H), 2.06-1.94 (m, 1H), 0.79 (d, J = 6.6 Hz, 3H), 0.76

- 53 -

(d, J = 6.6 Hz, 3H); MS (thermospray): m/z 374 (M+H).

EXAMPLE 18

2-(R)-2-[(2-tert-Butoxycarbonylethyl)(4-methoxybenzenesulfonyl)-amino]-N-hydroxy-3-methylbutyramide

A solution of N,N-dimethylformamide di-tert-butyl acetal (1.9 mL, 7.9 mmol) in toluene (15 mL) was added dropwise to a solution of 2-(R)-2-[(2-carboxyethyl(4-methoxybenzenesulfonyl)amino)-3-methylbutyric acid benzyl ester (example 2) 900 mg, 2.0 mmol) in toluene at 80°C. After heating for 2 hours at 80°C., the mixture was cooled and concentrated to leave an amber oil which was chromatographed on silica gel eluting with chloroform to afford 2-(R)-2-[(2-tert-butoxycarbonylethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester as an oil, 3.75 mg (37%).

To a solution of 2-(R)-2-[(2-tert-butoxycarbonylethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid benzyl ester (370 mg, 0.73 mmol) in ethanol (20 mL) was added 10% palladium on activated carbon (40 mg). The mixture was agitated under 3 atmospheres hydrogen in a Parr shaker for 5 hours. The catalyst was removed by filtration through diatomaceous earth and the solvent was evaporated leaving 2-(R)-2-[(2-tert-butoxycarbonylethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid as a white foam, 30 mg (100%).

To a solution of 2-(R)-2-[(2-tert-butoxycarbonylethyl)(4-methoxybenzenesulfonyl)amino]-3-methylbutyric acid (303 mg, 0.73 mmol) and 1-hydroxybenztriazole hydrate (108 mg, 0.70 mmol) in dry dimethylformamide (10 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (153 mg, 0.80 mmol). After stirring for 45 minutes, hydroxylamine hydrochloride (203 mg, 2.9 mmol) and then N-methylmorpholine (0.48 mL, 4.4 mmol) were added. The

- 54 -

mixture was stirred at room temperature overnight and then concentrated under vacuum. The residue was chromatographed on silica gel eluting with 2% methanol in chloroform and again with 10% ethyl acetate in hexane to afford 2-(R)-2-[(2-tert-butoxycarbonylethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide as a white foam, 135 mg (43%);  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  10.77 (br s, 1H), 7.74 (d,  $J = 8.9$  Hz, 2H), 7.08 (d,  $J = 8.9$  Hz, 2H), 3.93-3.82 (m, 1H, overlapped), 3.83 (s, 3H), 3.64 (d,  $J = 10.8$  Hz, 1H), 3.26-3.14 (m, 1H), 2.70-2.60 (m, 1H), 2.50-2.38 (m, 1H), 2.04-1.91 (m, 1H), 1.38 (s, 9H), 0.78 (d,  $J = 6.5$  Hz, 3H), 0.72 (d,  $J = 6.5$  Hz, 3H); MS (thermospray):  $m/z$  431 ( $M+H$ ), 375, 314.

15

EXAMPLE 19

2-(R)-2-[2-Carboxyethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide


To a solution of 2-(R)-2-[2-tert-butoxycarbonylethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide (example 18) (100 mg, 0.23 mmol) in methylene chloride (1 mL) at 0°C. was added trifluoroacetic acid (1 mL). The mixture was allowed to warm to room temperature while stirring overnight. After evaporation of the trifluoroacetic acid and methylene chloride, the residue was chromatographed on silica gel eluting with 5% methanol in chloroform. Concentration of the appropriate fractions afforded 2-(R)-2-[2-carboxyethyl)(4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide as a white solid, 35 mg (41%).  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ):  $\delta$  10.79 (br s, 1H), 8.97 (br s, 1H), 7.76 (d,  $J = 8.9$  Hz, 2H), 7.09 (d,  $J = 8.9$  Hz, 2H), 3.95-3.82 (m, 1H, overlapped), 3.84 (s, 3H), 3.66 (d,  $J = 10.8$  Hz, 1H), 3.30-3.20 (m, 1H), 2.73-2.62 (m, 1H), 2.50-2.40 (m, 1H), 2.07-1.94 (m, 1H), 0.80 (d,  $J = 6.5$  Hz, 3H), 0.74 (d,  $J = 6.5$  Hz, 3H); MS (thermospray):  $m/z$  375 ( $M+H$ ), 314.



- 55 -

C L A I M S

1. A compound of the formula




15 or the pharmaceutically acceptable salts thereof,  
wherein

n is 1 to 6;


X is hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or NR<sup>1</sup>R<sup>2</sup> wherein R<sup>1</sup> and R<sup>2</sup> are each independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, piperidyl, (C<sub>1</sub>-C<sub>6</sub>)alkylpiperidyl, (C<sub>6</sub>-C<sub>10</sub>)arylpiperidyl, (C<sub>5</sub>-C<sub>9</sub>)heteroarylpiperidyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperidyl, (C<sub>5</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperidyl, (C<sub>1</sub>-C<sub>6</sub>)acylpiperidyl, (C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>5</sub>-C<sub>9</sub>)heteroaryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>5</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>6</sub>-C<sub>10</sub>)aryl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sup>5</sup>(C<sub>2</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>5</sub>)alkyl(CHR<sup>5</sup>)(C<sub>1</sub>-C<sub>6</sub>)alkyl wherein R<sup>5</sup> is hydroxy, (C<sub>1</sub>-C<sub>6</sub>)acyloxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, piperazino, (C<sub>1</sub>-C<sub>6</sub>)acylamino, (C<sub>1</sub>-C<sub>6</sub>)alkylthio, (C<sub>6</sub>-C<sub>10</sub>)arylthio, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfinyl, (C<sub>6</sub>-C<sub>10</sub>)arylsulfinyl, (C<sub>1</sub>-C<sub>6</sub>)alkylsulfoxyl, (C<sub>6</sub>-C<sub>10</sub>)arylsulfoxyl, amino, (C<sub>1</sub>-C<sub>6</sub>)alkylamino, ((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>amino, (C<sub>1</sub>-C<sub>6</sub>)acylpiperazino, (C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino, (C<sub>5</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino, morpholino, thiomorpholino, piperidino or pyrrolidino; R<sup>6</sup>(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>5</sub>)alkyl(CHR<sup>6</sup>)(C<sub>1</sub>-C<sub>6</sub>)alkyl wherein R<sup>6</sup> is piperidyl, (C<sub>1</sub>-C<sub>6</sub>)alkylpiperidyl, (C<sub>6</sub>-C<sub>10</sub>)arylpiperidyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperidyl, (C<sub>5</sub>-

- $C_9$ )heteroarylpiperidyl or  $(C_5-C_9)$ heteroaryl $(C_1-C_6)$ alkylpiperidyl; and  $CH(R^7)COR^8$  wherein  $R^7$  is hydrogen,  $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkyl,  $(C_5-C_9)$ heteroaryl $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylthio $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ arylthio $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylsulfinyl $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ arylsulfinyl $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylsulfonyl $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ arylsulfonyl $(C_1-C_6)$ alkyl, hydroxy $(C_1-C_6)$ alkyl, amino $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylamino $(C_1-C_6)$ alkyl,  $((C_1-C_6)$ alkylamino) $_2$  $(C_1-C_6)$ alkyl,  $R^9R^{10}NCO(C_1-C_6)$ alkyl or  $R^9OCO(C_1-C_6)$ alkyl wherein  $R^9$  and  $R^{10}$  are each independently selected from the group consisting of hydrogen,  $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkyl and  $(C_5-C_9)$ heteroaryl $(C_1-C_6)$ alkyl; and  $R^8$  is  $R^{11}O$  or  $R^{11}R^{12}N$  wherein  $R^{11}$  and  $R^{12}$  are each independently selected from the group consisting of hydrogen,  $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkyl and  $(C_5-C_9)$ heteroaryl $(C_1-C_6)$ alkyl; or  $R^1$  and  $R^2$ , or  $R^9$  and  $R^{10}$ , or  $R^{11}$  and  $R^{12}$  may be taken together to form an azetidiny, pyrrolidinyl, morpholinyl, thiomorpholinyl, indolinyl, isoindolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl,  $(C_1-C_6)$ acylpiperazinyl,  $(C_1-C_6)$ alkylpiperazinyl,  $(C_6-C_{10})$ arylpiperazinyl,  $(C_5-C_9)$ heteroarylpiperazinyl or a bridged diazabicycloalkyl ring selected from the group consisting of




- 57 -

5



d



e

wherein  $r$  is 1, 2 or 3;

$m$  is 1 or 2;

10

$p$  is 0 or 1; and

$Q$  is hydrogen,  $(C_1-C_3)$ alkyl or  $(C_1-C_6)$ acyl;

$R^3$  and  $R^4$  are each independently selected

from the group consisting of hydrogen,  $(C_1-C_6)$ alkyl,

trifluoromethyl, trifluoromethyl $(C_1-C_6)$ alkyl,  $(C_1-$

15

$C_6)$ alkyl (difluoromethylene),  $(C_1-$

$C_3)$ alkyl (difluoromethylene)  $(C_1-C_3)$ alkyl,  $(C_6-C_{10})$ aryl,

$(C_5-C_9)$ heteroaryl,  $(C_6-C_{10})$ aryl  $(C_1-C_6)$ alkyl,  $(C_5-$

$C_9)$ heteroaryl  $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl  $(C_6-C_{10})$ aryl,

$(C_6-C_{10})$ aryl  $(C_6-C_{10})$ aryl  $(C_1-C_6)$ alkyl,  $(C_3-$

20

$C_6)$ cycloalkyl,  $(C_3-C_6)$ cycloalkyl  $(C_1-C_6)$ alkyl,

hydroxy $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ acyloxy $(C_1-C_6)$ alkyl,  $(C_1-$

$C_6)$ alkoxy $(C_1-C_6)$ alkyl, piperazinyl  $(C_1-C_6)$ alkyl,  $(C_1-$

$C_6)$ acylamino $(C_1-C_6)$ alkyl, piperidyl,  $(C_1-$

$C_6)$ alkylpiperidyl,  $(C_6-C_{10})$ aryl  $(C_1-C_6)$ alkoxy $(C_1-$

25

$C_6)$ alkyl,  $(C_5-C_9)$ heteroaryl  $(C_1-C_6)$ alkoxy $(C_1-C_6)$ alkyl,

$(C_1-C_6)$ alkylthio $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ arylthio $(C_1-$

$C_6)$ alkyl,  $(C_1-C_6)$ alkylsulfinyl  $(C_1-C_6)$ alkyl,  $(C_6-$

$C_{10})$ arylsulfinyl  $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylsulfonyl  $(C_1-$

$C_6)$ alkyl,  $(C_6-C_{10})$ arylsulfonyl  $(C_1-C_6)$ alkyl, amino $(C_1-$

30

$C_6)$ alkyl,  $(C_1-C_6)$ alkylamino $(C_1-C_6)$ alkyl,  $((C_1-$

$C_6)$ alkylamino) $_2$   $(C_1-C_6)$ alkyl,  $R^{13}CO(C_1-C_6)$ alkyl wherein

$R^{13}$  is  $R^{20}O$  or  $R^{20}R^{21}N$  wherein  $R^{20}$  and  $R^{21}$  are each

independently selected from the group consisting of

hydrogen,  $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl  $(C_1-C_6)$ alkyl or

35


$(C_5-C_9)$ heteroaryl  $(C_1-C_6)$ alkyl; or  $R^{14}(C_1-C_6)$ alkyl

wherein  $R^{14}$  is  $(C_1-C_6)$ acylpiperazino,  $(C_6-$

- 58 -

- $C_{10}$ )arylpiperazino,  $(C_5-C_9)$ heteroarylpiperazino,  $(C_1-C_6)$ alkylpiperazino,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkylpiperazino,  $(C_5-C_9)$ heteroaryl $(C_1-C_6)$ alkylpiperazino, morpholino, thiomorpholino, piperidino, pyrrolidino, piperidyl,
- 5  $(C_1-C_6)$ alkylpiperidyl,  $(C_6-C_{10})$ arylpiperidyl,  $(C_5-C_9)$ heteroarylpiperidyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkylpiperidyl,  $(C_5-C_9)$ heteroaryl $(C_1-C_6)$ alkylpiperidyl or  $(C_1-C_6)$ acylpiperidyl;
- or  $R^3$  and  $R^4$ , or  $R^{20}$  and  $R^{21}$  may be taken
- 10 together to form a  $(C_3-C_6)$ cycloalkyl, oxacyclohexyl, thiocyclohexyl, indanyl or tetralinyl ring or a group of the formula

15



- wherein  $R^{15}$  is hydrogen,  $(C_1-C_6)$ acyl,  $(C_1-C_6)$ alkyl,
- 20  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkyl,  $(C_5-C_9)$ heteroaryl $(C_1-C_6)$ alkyl or  $(C_1-C_6)$ alkylsulfonyl; and

- Ar is  $(C_6-C_{10})$ aryl,  $(C_5-C_9)$ heteroaryl,  $(C_1-C_6)$ alkyl $(C_6-C_{10})$ aryl,  $(C_1-C_6)$ alkoxy $(C_6-C_{10})$ aryl,  $((C_1-C_6)$ alkoxy) $_2$  $(C_6-C_{10})$ aryl,  $(C_6-C_{10})$ aryloxy $(C_6-C_{10})$ aryl,
- 25  $(C_5-C_9)$ heteroaryloxy $(C_6-C_{10})$ aryl,  $(C_1-C_6)$ alkyl $(C_5-C_9)$ heteroaryl,  $(C_1-C_6)$ alkoxy $(C_5-C_9)$ heteroaryl,  $((C_1-C_6)$ alkoxy) $_2$  $(C_5-C_9)$ heteroaryl,  $(C_6-C_{10})$ aryloxy $(C_5-C_9)$ heteroaryl,  $(C_5-C_9)$ heteroaryloxy $(C_5-C_9)$ heteroaryl;

- with the proviso that when either  $R^1$  or  $R^2$  is
- 30  $CH(R^7)COR^8$  wherein  $R^7$  and  $R^8$  are as defined above, the other of  $R^1$  or  $R^2$  is hydrogen,  $(C_1-C_6)$ alkyl or benzyl.

2. A compound according to claim 1, wherein n is 2.
3. A compound according to claim 1, wherein Ar
- 35 is 4-methoxyphenyl or 4-phenoxyphenyl.
4. A compound according to claim 1, 2 or 3, wherein either  $R^3$  or  $R^4$  is not hydrogen.
5. A compound according to claim 1, wherein n is


- 59 -

1 and either  $R^1$  or  $R^2$  is hydrogen.

6. A compound according to claim 4, wherein X is hydroxy, Ar is 4-methoxyphenyl or 4-phenoxyphenyl.

7. A compound according to claim 4, wherein X is alkoxy, Ar is 4-methoxyphenyl or 4-phenoxyphenyl.

8. A compound according to claim 1, wherein Ar is 4-methoxyphenyl or 4-phenoxyphenyl and  $R^3$  and  $R^4$  are taken together to form (C<sub>3</sub>-C<sub>6</sub>)cycloalkanyl, oxacyclohexanyl, thiocyclohexanyl, indanyl or a group of the formula



wherein  $R^{15}$  is (C<sub>1</sub>-C<sub>6</sub>)acyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>5</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>1</sub>-C<sub>6</sub>)alkylsulfonyl.

9. A compound according to claim 1, wherein n is 2, Ar is 4-methoxyphenyl or 4-phenoxyphenyl,  $R^1$  and  $R^2$  are taken together to form piperazinyl, (C<sub>1</sub>-C<sub>6</sub>)alkylpiperazinyl, (C<sub>6</sub>-C<sub>10</sub>)aryl piperazinyl or (C<sub>5</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperazinyl, and either  $R^3$  or  $R^4$  is not hydrogen or both  $R^3$  and  $R^4$  are not hydrogen.

10. A compound according to claim 1, wherein n is 2, Ar is 4-methoxyphenyl or 4-phenoxyphenyl,  $R^1$  is hydrogen or (C<sub>1</sub>-C<sub>6</sub>)alkyl,  $R^2$  is 2-pyridylmethyl, 3-pyridylmethyl or 4-pyridylmethyl, and either  $R^3$  or  $R^4$  is not hydrogen or both  $R^3$  and  $R^4$  are not hydrogen.

11. A compound according to claim 1, wherein n is 1, Ar is 4-methoxyphenyl or 4-phenoxyphenyl,  $R^1$  is hydrogen,  $R^2$  is 2-pyridylmethyl, 3-pyridylmethyl or 4-pyridylmethyl, and either  $R^3$  or  $R^4$  is not hydrogen or both  $R^3$  and  $R^4$  are not hydrogen.

12. A compound according to claim 2, wherein Ar is 4-methoxyphenyl,  $R^1$  is hydrogen or (C<sub>1</sub>-C<sub>6</sub>)alkyl and  $R^2$  is  $R^5$ (C<sub>2</sub>-C<sub>6</sub>)alkyl wherein  $R^5$  is morpholino,

thiomorpholino, piperidino, pyrrolidino, (C<sub>1</sub>-C<sub>6</sub>)acylpiperazino, (C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino, (C<sub>6</sub>-C<sub>10</sub>)arylpiperazino, (C<sub>5</sub>-C<sub>9</sub>)heteroarylpiperazino, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino or (C<sub>5</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino and either R<sup>3</sup> or R<sup>4</sup> is not hydrogen or both R<sup>3</sup> and R<sup>4</sup> are not hydrogen.

13. A compound according to claim 1, wherein n is 1, Ar is 4-methoxyphenyl or 4-phenoxyphenyl, R<sup>1</sup> is hydrogen, R<sup>2</sup> is R<sup>5</sup>(C<sub>2</sub>-C<sub>6</sub>)alkyl wherein R<sup>5</sup> is morpholino, thiomorpholino, piperidino, pyrrolidino, (C<sub>1</sub>-C<sub>6</sub>)acylpiperazino, (C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino, (C<sub>6</sub>-C<sub>10</sub>)arylpiperazino, (C<sub>5</sub>-C<sub>9</sub>)heteroarylpiperazino, (C<sub>6</sub>-C<sub>10</sub>)aryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino or (C<sub>5</sub>-C<sub>9</sub>)heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkylpiperazino and either R<sup>3</sup> or R<sup>4</sup> is not hydrogen or both R<sup>3</sup> and R<sup>4</sup> are not hydrogen.

14. A compound according to claim 1, wherein said compound is selected from:

- 2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)(3-morpholin-4-yl-3-oxopropyl)amino]-3-methylbutyramide;
- 2-(R)-2-[(2-Benzylcarbamoyl)ethyl](4-methoxybenzenesulfonyl)amino]-N-hydroxy-3-methylbutyramide;
- 2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)(2-[(pyridin-3-ylmethyl)carbamoyl]ethyl)amino]-3-methylbutyramide;
- 2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)[2-(methylpyridin-3-ylmethylcarbamoyl)ethyl]amino]-3-methylbutyramide;
- 4-(3-[1-(R)-1-Hydroxycarbamoyl-2-methylpropyl](4-methoxybenzenesulfonyl)amino)propionyl)piperazine-1-carboxylic acid, tert-butyl ester;
- 2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)(3-oxo-3-piperazin-1-ylpropyl)amino]-3-methylbutyramide hydrochloride;
- 2-(R)-2-[(Benzylcarbamoylmethyl)(4-

- 61 -

- methoxybenzenesulfonyl) amino] N-hydroxy-3-methylbutyramide;
- 2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)-(2-morpholin-4-ylethylcarbamoyl)methyl] amino)-3-methylbutyramide;
- 5 2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)-[(pyridin-3-ylmethyl)carbamoyl)methyl] amino)-3-methylbutyramide;
- 2-(R)-3,3,3-Trifluoro-N-hydroxy-2-[(methoxybenzenesulfonyl)(3-morpholin-4-yl-3-oxopropyl) amino] propionamide;
- 10 2-(R)-N-Hydroxy-2-[(4-phenoxybenzenesulfonyl)[2-(methylpyridin-4-ylmethylcarbamoyl) ether] amino)-3-methylbutyramide;
- 15 4-[4-Methoxybenzenesulfonyl)(3-morpholin-4-yl-3-oxopropyl) amino]-1-methylpiperidene-4-carboxylic acid hydroxyamide;
- 2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)-[3-(4-methylpiperazin-1-yl)-3-oxopropyl] amino)-3-methylbutyramide;
- 20 2-(R)-2-[(2-Carboxyethyl)(4-methoxybenzenesulfonyl) amino]-N-hydroxy-3-methylbutyramide;
- [(2-Carboxyethyl)(3,4-dimethoxybenzenesulfonyl) amino]-N-hydroxy-acetamide;
- 25 2-(R)-2-[(2-Carbamoylethyl)(4-methoxybenzenesulfonyl) amino]-N-hydroxy-3-methylbutyramide;
- 2-(R),3-(R)-3,N-Dihydroxy-2-[(4-methoxybenzenesulfonyl)(3-oxo-3-piperidin-1-ylpropyl) amino]-butyramide;
- 30 2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)[3-(methylpyridin-3-ylmethylcarbamoyl)propyl] amino)-3-methylbutyramide;
- 35 2-(R)-N-Hydroxy-2-[(4-methoxybenzenesulfonyl)[2-(methylcarboxymethylcarbamoyl)ethyl] amino)-3-

- 62 -

methylbutyramide;

2-(R)-N-Hydroxy-2-((4-methoxybenzenesulfonyl)-[(1-methylpiperidin-4-ylcarbonyl)methyl]amino)-3-methylbutyramide;

5 2-(R)-2-Cyclohexyl-N-hydroxy-2-((4-methoxybenzenesulfonyl)-[3-(4-methylpiperazin-1-yl)-3-oxopropyl]amino)-acetamide; and

2-(R)-N-Hydroxy-2-[(methoxybenzenesulfonyl)(3-morpholin-4-yl-3-oxopropyl)amino]-4-(morpholin-4-yl)butyramide.

10 15. A pharmaceutical composition for (a) the treatment of a condition selected from the group consisting of arthritis, cancer, tissue ulceration, restenosis, periodontal disease, epidermolysis bullosa, scleritis and other diseases characterized by matrix metalloproteinase activity, AIDS, sepsis, septic shock and other diseases involving the production of tumor necrosis factor (TNF) or (b) the inhibition of matrix metalloproteinases or the production of tumor necrosis factor (TNF) in a mammal, including a human, comprising an amount of a compound of claim 1 or a pharmaceutically acceptable salt thereof, effective in such treatments and a pharmaceutically acceptable carrier.

25 16. A method for the inhibition of (a) matrix metalloproteinases or (b) the production of tumor necrosis factor (TNF) in a mammal, including a human, comprising administering to said mammal an effective amount of a compound of claim 1 or a pharmaceutically acceptable salt thereof.


30 17. A method for treating a condition selected from the group consisting of arthritis, cancer, tissue ulceration, restenosis, periodontal disease, epidermolysis bullosa, scleritis and other diseases characterized by matrix metalloproteinase activity, AIDS, sepsis, septic shock and other diseases involving the production of tumor necrosis factor (TNF) in a



- 63 -

mammal, including a human, comprising administering to said mammal an amount of a compound of claim 1 or a pharmaceutically acceptable salt thereof, effective in treating such a condition.

- 5 18. A method of preparing a compound of the formula



- 15 or the pharmaceutically acceptable salts thereof, wherein

$n$  is 1 to 6;


- $X$  is hydroxy,  $(C_1-C_6)$ alkoxy or  $NR^1R^2$  wherein  $R^1$  and  $R^2$  are each independently selected from the group consisting of hydrogen,  $(C_1-C_6)$ alkyl, piperidyl,  $(C_1-C_6)$ alkylpiperidyl,  $(C_6-C_{10})$ aryl piperidyl,  $(C_5-C_9)$ heteroaryl piperidyl,  $(C_6-C_{10})$ aryl  $(C_1-C_6)$ alkylpiperidyl,  $(C_5-C_9)$ heteroaryl  $(C_1-C_6)$ alkylpiperidyl,  $(C_1-C_6)$ acylpiperidyl,  $(C_6-C_{10})$ aryl,  $(C_5-C_9)$ heteroaryl,  $(C_6-C_{10})$ aryl  $(C_1-C_6)$ alkyl,  $(C_5-C_9)$ heteroaryl  $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl  $(C_6-C_{10})$ aryl,  $(C_6-C_{10})$ aryl  $(C_6-C_{10})$ aryl  $(C_1-C_6)$ alkyl,  $(C_3-C_6)$ cycloalkyl,  $(C_3-C_6)$ cycloalkyl  $(C_1-C_6)$ alkyl,  $R^5(C_2-C_6)$ alkyl,  $(C_1-C_5)$ alkyl  $(CHR^5)$   $(C_1-C_6)$ alkyl wherein  $R^5$  is hydroxy,  $(C_1-C_6)$ acyloxy,  $(C_1-C_6)$ alkoxy, piperazino,  $(C_1-C_6)$ acylamino,  $(C_1-C_6)$ alkylthio,  $(C_6-C_{10})$ arylthio,  $(C_1-C_6)$ alkylsulfinyl,  $(C_6-C_{10})$ arylsulfinyl,  $(C_1-C_6)$ alkylsulfoxyl,  $(C_6-C_{10})$ arylsulfoxyl, amino,  $(C_1-C_6)$ alkylamino,  $((C_1-C_6)alkyl)_2$ amino,  $(C_1-C_6)$ acylpiperazino,  $(C_1-C_6)$ alkylpiperazino,  $(C_6-C_{10})$ aryl  $(C_1-C_6)$ alkylpiperazino,  $(C_5-C_9)$ heteroaryl  $(C_1-$
- 20
- 25
- 30
- 35

- 64 -


- $C_6$ )alkylpiperazino, morpholino, thiomorpholino, piperidino or pyrrolidino;  $R^6$  ( $C_1-C_6$ )alkyl, ( $C_1-C_5$ )alkyl( $CHR^6$ ) ( $C_1-C_6$ )alkyl wherein  $R^6$  is piperidyl, ( $C_1-C_6$ )alkylpiperidyl, ( $C_6-C_{10}$ )arylpiperidyl, ( $C_6-C_{10}$ )aryl( $C_1-C_6$ )alkylpiperidyl, ( $C_5-C_9$ )heteroarylpiperidyl or ( $C_5-C_9$ )heteroaryl( $C_1-C_6$ )alkylpiperidyl; and  $CH(R^7)COR^8$  wherein  $R^7$  is hydrogen, ( $C_1-C_6$ )alkyl, ( $C_6-C_{10}$ )aryl( $C_1-C_6$ )alkyl, ( $C_5-C_9$ )heteroaryl( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkylthio( $C_1-C_6$ )alkyl, ( $C_6-C_{10}$ )arylthio( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkylsulfinyl( $C_1-C_6$ )alkyl, ( $C_6-C_{10}$ )arylsulfinyl( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkylsulfonyl( $C_1-C_6$ )alkyl, ( $C_6-C_{10}$ )arylsulfonyl( $C_1-C_6$ )alkyl, hydroxy( $C_1-C_6$ )alkyl, amino( $C_1-C_6$ )alkyl, ( $C_1-C_6$ )alkylamino( $C_1-C_6$ )alkyl, (( $C_1-C_6$ )alkylamino)<sub>2</sub>( $C_1-C_6$ )alkyl,  $R^9R^{10}NCO(C_1-C_6)alkyl$  or  $R^9OCO(C_1-C_6)alkyl$  wherein  $R^9$  and  $R^{10}$  are each independently selected from the group consisting of hydrogen, ( $C_1-C_6$ )alkyl, ( $C_6-C_{10}$ )aryl( $C_1-C_6$ )alkyl and ( $C_5-C_9$ )heteroaryl( $C_1-C_6$ )alkyl; and  $R^8$  is  $R^{11}O$  or  $R^{11}R^{12}N$  wherein  $R^{11}$  and  $R^{12}$  are each independently selected from the group consisting of hydrogen, ( $C_1-C_6$ )alkyl, ( $C_6-C_{10}$ )aryl( $C_1-C_6$ )alkyl and ( $C_5-C_9$ )heteroaryl( $C_1-C_6$ )alkyl;
- or  $R^1$  and  $R^2$ , or  $R^9$  and  $R^{10}$ , or  $R^{11}$  and  $R^{12}$
- may be taken together to form an azetidiny, pyrrolidinyl, morpholinyl, thiomorpholinyl, indolinyl, isoindolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, ( $C_1-C_6$ )acylpiperazinyl, ( $C_1-C_6$ )alkylpiperazinyl, ( $C_6-C_{10}$ )arylpiperazinyl, ( $C_5-C_9$ )heteroarylpiperazinyl or a bridged diazabicycloalkyl ring selected from the group consisting of

- 65 -


5



a




b




c

10



d



e

wherein  $r$  is 1, 2 or 3;

15

$m$  is 1 or 2;

$p$  is 0 or 1; and

$Q$  is hydrogen,  $(C_1-C_3)$ alkyl or  $(C_1-C_6)$ acyl;

$R^3$  and  $R^4$  are each independently selected

20

from the group consisting of hydrogen,  $(C_1-C_6)$ alkyl,

trifluoromethyl, trifluoromethyl $(C_1-C_6)$ alkyl,  $(C_1-$

$C_6)$ alkyl (difluoromethylene),  $(C_1-$

$C_3)$ alkyl (difluoromethylene)  $(C_1-C_3)$ alkyl,  $(C_6-C_{10})$ aryl,

$(C_5-C_9)$ heteroaryl,  $(C_6-C_{10})$ aryl  $(C_1-C_6)$ alkyl,  $(C_5-$

$C_9)$ heteroaryl  $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl  $(C_6-C_{10})$ aryl,

25

$(C_6-C_{10})$ aryl  $(C_6-C_{10})$ aryl  $(C_1-C_6)$ alkyl,  $(C_3-$

$C_6)$ cycloalkyl,  $(C_3-C_6)$ cycloalkyl  $(C_1-C_6)$ alkyl,

hydroxy  $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ acyloxy  $(C_1-C_6)$ alkyl,  $(C_1-$

$C_6)$ alkoxy  $(C_1-C_6)$ alkyl, piperazinyl  $(C_1-C_6)$ alkyl,  $(C_1-$

$C_6)$ acylamino  $(C_1-C_6)$ alkyl, piperidyl,  $(C_1-$

30

$C_6)$ alkylpiperidyl,  $(C_6-C_{10})$ aryl  $(C_1-C_6)$ alkoxy  $(C_1-$


$C_6)$ alkyl,  $C_5-C_9)$ heteroaryl  $(C_1-C_6)$ alkoxy  $(C_1-C_6)$ alkyl,

$(C_1-C_6)$ alkylthio  $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ arylthio  $(C_1-$

- 66 -

$C_6$ )alkyl,  $(C_1-C_6)$ alkylsulfinyl $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ arylsulfinyl $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylsulfonyl $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ arylsulfonyl $(C_1-C_6)$ alkyl, amino $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkylamino $(C_1-C_6)$ alkyl,  $((C_1-C_6)$ alkylamino) $_2$  $(C_1-C_6)$ alkyl,  $R^{13}CO(C_1-C_6)$ alkyl wherein  $R^{13}$  is  $R^{20}O$  or  $R^{20}R^{21}N$  wherein  $R^{20}$  and  $R^{21}$  are each independently selected from the group consisting of hydrogen,  $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkyl or  $(C_5-C_9)$ heteroaryl $(C_1-C_6)$ alkyl; or  $R^{14}(C_1-C_6)$ alkyl wherein  $R^{14}$  is  $(C_1-C_6)$ acylpiperazino,  $(C_6-C_{10})$ arylpiperazino,  $(C_5-C_9)$ heteroarylpiperazino,  $(C_1-C_6)$ alkylpiperazino,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkylpiperazino,  $(C_5-C_9)$ heteroaryl $(C_1-C_6)$ alkylpiperazino, morpholino, thiomorpholino, piperidino, pyrrolidino, piperidyl,  $(C_1-C_6)$ alkylpiperidyl,  $(C_6-C_{10})$ arylpiperidyl,  $(C_5-C_9)$ heteroarylpiperidyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkylpiperidyl,  $(C_5-C_9)$ heteroaryl $(C_1-C_6)$ alkylpiperidyl or  $(C_1-C_6)$ acylpiperidyl; or  $R^3$  and  $R^4$ , or  $R^{20}$  and  $R^{21}$  may be taken together to form a  $(C_3-C_6)$ cycloalkyl, oxacyclohexyl, thiocyclohexyl, indanyl or tetralinyl ring or a group of the formula


25



wherein  $R^{15}$  is hydrogen,  $(C_1-C_6)$ acyl,  $(C_1-C_6)$ alkyl,  $(C_6-C_{10})$ aryl $(C_1-C_6)$ alkyl,  $(C_5-C_9)$ heteroaryl $(C_1-C_6)$ alkyl or  $(C_1-C_6)$ alkylsulfonyl; and  
 Ar is  $(C_6-C_{10})$ aryl,  $(C_5-C_9)$ heteroaryl,  $(C_1-C_6)$ alkyl $(C_6-C_{10})$ aryl,  $(C_1-C_6)$ alkoxy $(C_6-C_{10})$ aryl,  $((C_1-C_6)$ alkoxy) $_2$  $(C_6-C_{10})$ aryl,  $(C_6-C_{10})$ aryloxy $(C_6-C_{10})$ aryl,  $(C_5-C_9)$ heteroaryloxy $(C_6-C_{10})$ aryl,  $(C_1-C_6)$ alkyl $(C_5-C_9)$ heteroaryl,  $(C_1-C_6)$ alkoxy $(C_5-C_9)$ heteroaryl,  $((C_1-C_6)$ alkoxy) $_2$  $(C_5-C_9)$ heteroaryl,  $(C_6-C_{10})$ aryloxy $(C_5-C_9)$ heteroaryl,  $(C_5-C_9)$ heteroaryloxy $(C_5-C_9)$ heteroaryl;

- 67 -

with the proviso that when either  $R^1$  or  $R^2$  is  $CH(R^7)COR^8$  wherein  $R^7$  and  $R^8$  are as defined above, the other of  $R^1$  or  $R^2$  is hydrogen,  $(C_1-C_6)$ alkyl or benzyl; comprising reacting a compound of the formula



wherein  $n$ ,  $X$ ,  $R^3$ ,  $R^4$  and  $\text{Ar}$  are as defined above with  
 1- (3-dimethylaminopropyl) -3-ethylcarbodiimide, 1-  
 15 hydroxybenztriazole and hydroxylamine.

# INTERNATIONAL SEARCH REPORT

In:      onal Application No  
PCT/US 96/02679

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6    C07C311/29    C07D295/18    C07D213/56    A61K31/535    A61K31/44  
         A61K31/495    A61K31/40    A61K31/18

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6    C07C    C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages       | Relevant to claim No. |
|------------|--|-----------------------|
| X          | EP,A,0 606 046 (CIBA-GEIGY AG) 13 July 1994<br>see the whole document<br>---             | 1-15,18               |
| A          | WO,A,90 05719 (BRITISH BIOTECHNOLOGY LTD) 31 May 1990<br>see the whole document<br>----- | 1-15,18               |

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- \*&\* document member of the same patent family

Date of the actual completion of the international search

15 May 1996

Date of mailing of the international search report

04.06.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+ 31-70) 340-3016

Authorized officer

Goetz, G

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No  
PCT/US 96/02679

| Patent document<br>cited in search report | Publication<br>date | Patent family<br>member(s) | Publication<br>date |
|---|---------------------|----------------------------|---------------------|
| EP-A-606046                               | 13-07-94            | US-A- 5455258              | 03-10-95            |
|   |                     | AU-B- 5265593              | 04-05-95            |
|   |                     | CA-A- 2112779              | 07-07-94            |
|   |                     | FI-A- 940012               | 07-07-94            |
|   |                     | HU-A- 70536                | 30-10-95            |
|   |                     | JP-A- 6256293              | 13-09-94            |
|   |                     | NO-A- 940038               | 07-07-94            |
|   |                     | NZ-A- 250517               | 26-10-95            |
|   |                     | US-A- 5506242              | 09-04-96            |
|   |                     | ZA-A- 9400048              | 11-08-94            |
| -----                                     |                     |                            |                     |
| WO-A-9005719                              | 31-05-90            | AU-B- 644064               | 02-12-93            |
|   |                     | AU-B- 4800390              | 12-06-90            |
|   |                     | CA-A- 2003718              | 23-05-90            |
|   |                     | DE-D- 68914687             | 19-05-94            |
|   |                     | DE-T- 68914687             | 08-09-94            |
|   |                     | EP-A- 0446267              | 18-09-91            |
|   |                     | ES-T- 2055409              | 16-08-94            |
|   |                     | JP-T- 4502008              | 09-04-92            |
|   |                     | NO-B- 177701               | 31-07-95            |
|   |                     | US-A- 5310763              | 10-05-94            |
| US-A- 5240958                             | 31-08-93            |                            |                     |
| -----                                     |                     |                            |                     |